

# A Survey on Relationship between Diet and Urinary Excretion of Aflatoxin M<sub>1</sub>: A Screening Pilot Study on Iranian Population

S. Mason<sup>1</sup>, B. Hajimohammadi<sup>2</sup>, M.H. Ehrampoush<sup>2</sup>, F. Khabiri<sup>2</sup>, M. Soltani<sup>2\*</sup>

1. School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, UK

2. Research Center for Molecular Identification of Food Hazards, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

## Article type

Original article

## Abstract

### Keywords

Aflatoxins  
Biological Markers  
Food Safety  
Iran

Received: 1 Dec 2014

Revised: 29 Dec 2014

Accepted: 18 Jan 2015

**Background:** Aflatoxins are found in the most types of foods such as corn, peanuts, pistachio, rice, milk, and etc. These toxins may cause several adverse health effects, notably cancer. The main aim of this study was to evaluate excretion level of aflatoxin M<sub>1</sub> (AFM<sub>1</sub>), as a biomarker of aflatoxin B<sub>1</sub> exposure, in urine samples of Iranian population. Also, relationship between diet and urinary excretion of AFM<sub>1</sub> has been discussed.

**Methods:** From June to August 2014, this study was carried out among 70 staffs from Faculty of Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. From each participant, 72-h dietary recall was asked and recorded. Then, urine samples were collected separately in the sterile plastic falcon and transferred immediately to the laboratory and stored at -20 °C. AFM<sub>1</sub> was assessed in each sample using ELISA procedure. The data were analyzed by Kruskal–Wallis one-way analysis of variance test using SPSS software (16.0).

**Results:** AFM<sub>1</sub> was detected in 15 of 70 (21%) urine samples. The mean levels of contamination in faculty members and non-faculty members were 0.6 and 1.7 pg/ml, respectively. There was a significant difference between the excretion of AFM<sub>1</sub> and consumption of nuts as well as Iranian traditional confection ( $p < 0.05$ ). No relationship was found between the demographic factors and excretion of AFM<sub>1</sub> ( $p > 0.05$ ).

**Conclusions:** Considering adverse health effect, aflatoxin exposure in this Iranian population should be reduced. So, comprehensive training of the people is one of the most practical and useful methods for reducing aflatoxin health risks.

## Introduction

Aflatoxin (AF) is a group of mycotoxin compounds, which is produced in foods by three main species of *Aspergillus* genus including *A. flavus*, *A. parasiticus* and *A. nomius* (Atanda et al., 2007; Ehrlich et al., 2007; Khodadadi et al., 2014). AFs are mainly found in rice (Feizy et al., 2010), sesame (Asadi et al., 2011), nuts (Pour et al., 2010), and milk as well as dairy products (Rahimi et al., 2010).

Typically, there are four types of AFs including AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub> that are recognized as high toxic compounds. Liver is the main organ affected by these carcinogenic toxins (Groopman and Kensler, 2005). The existence of AFs in foods may cause chronic health effects including immune-system suppression (Shuaib et al., 2010), impaired childhood development (Sherif et al., 2009) and notably cancer. Also, acute toxicity can lead to

\*Corresponding author

E-mail: soltani.mansooreh@gmail.com

death (Iha et al., 2011; Smith et al., 2012; Yazdanpanah et al., 2013).

Exposure assessment is one of the most important parts of the risk assessment process (Sherif et al., 2009; Wu et al., 2009). Dietary intake of AF contaminated food is the main source of human exposure to AFs (Egal et al., 2005; Lewis et al., 2005). Human exposure to AFB<sub>1</sub> has been determined by excretion of its metabolites in urine, milk and blood. Approximately, 1.2-2.2% of the AF received through food is excreted as AFM<sub>1</sub> in urine of human (Asadi et al., 2011; Cheraghali et al., 2007; Leong et al., 2012; Mahdavi et al., 2010; Pour et al., 2010; Rahimi et al., 2010; Romero et al., 2010; Sadeghi et al., 2009; Zaghini et al., 2005).

The presence of AFM<sub>1</sub>, a hydroxylated metabolite of AFB<sub>1</sub> and one of the most important AF metabolites, in breast milk provides main source of its exposure for lactating newborn infants and breast feeding (Afshar et al., 2013; Battacone et al., 2005; Iha et al., 2011; Leong et al., 2012; Mahdavi et al., 2010; Mckean et al., 2006; Partanen et al., 2009; Polychronaki et al., 2008; Sadeghi et al., 2009; Yazdanpanah et al., 2013; Zaghini et al., 2005).

Among some previous similar published studies, Sabran et al. (2012) found significant relationship between consumption of milk and dairy products and excretion of AFM<sub>1</sub>. Romero et al. (2010) evaluated excretion of AFM<sub>1</sub> in 69 people in Brazil, 45 of them (65%) had contamination. Polychronaki et al. (2008) detected AFM<sub>1</sub> in 38% of Egyptian and also 86% of Guinean urine samples.

As it was thought the staffs of Faculty of Health had enough knowledge of AF and its risks; it was expected that they may have less exposure to it in comparison with others. Therefore, in this study, we decided to investigate relationship between diet and urinary excretion of AFM<sub>1</sub> in staffs of Faculty of Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

## Material and methods

### Ethical approval

This research was approved by Shahid Sadoughi University of Medical Sciences, Yazd, Iran (Protocol No. 3493).

Informed consent was obtained from each participant before beginning the study.

### Sampling

This study was a cross sectional-descriptive study that was done from June to August 2014. Seventy people were selected among the eligible employees including academic and non-academic staffs.

Urine samples were collected separately in the plastic

falcons and transferred immediately to the laboratory and stored at -20 °C.

### Determination of the AFM<sub>1</sub>

ELISA kit (6827 BN Arnhem, Euro Proxima Company, Arnhem, Netherlands) was used for detection of AFM<sub>1</sub>. The sample preparation was performed according to the manufacturer's instructions. All samples were centrifuged for 10 min in 2000 ×g at 4 °C. Each urine sample was diluted 1:1 with sample dilution buffer, mixed well and distributed in 100 µl inside the respective wells of the plate.

According to the manufacturer's instructions, the AFM<sub>1</sub> standards and test urine samples were added in duplicate to each well plate, and then incubated at room temperature (20-25 °C) for 1 h in darkness. After that, the solution was discarded from the microtitre plate and washed 3 times with rinsing buffer. Then, 100 µl of conjugate (Aflatoxin M<sub>1</sub> -HRPO) were added to each well, except zero standard maximal wells. The microtitre plate was shaken for few seconds on a microtitre plate shaker, and then incubated at room temperature for 30 min in darkness. The solution was discarded from the microtitre plate and washed 3 times with rinsing buffer. Amount of 100 µl of substrate solution were added into each well and incubated at room temperature for 30 min.

Finally, 100 µl of stop solution were added to each well. The optical absorbance was read at 450 nm with microplate reader (ELX 800 UV, Bio-Tek Instruments, Inc).

### Statistical analysis

Kruskal-Wallis one-way analysis of variance test was used to determine the relationship between type of dairy products, type of rice, place of residence and demographic factors with the concentration of AFM<sub>1</sub>. Mann-Whitney test was used to compare the two groups and determine the relationship between food intake and the concentration of AFM<sub>1</sub> in urine. The data were analyzed using SPSS software (16.0).

## Results

AFM<sub>1</sub> was detected in 15 of 70 (21%) urine samples of the studied population (Table 1). The mean levels of contamination in academic and non-academic staffs were 0.6 and 1.7 pg/ml, respectively.

Results of this study revealed that no significant relationship was observed between academic and non-academic staffs with excretion level of AFM<sub>1</sub> ( $p > 0.05$ ). There was a significant difference ( $p < 0.05$ ) between the excretion level of AFM<sub>1</sub> and consumption of nuts as well as Iranian traditional confection (Table 2). No relation-

ship ( $p > 0.05$ ) was found between the socio-demographic factors and excretion level of AFM<sub>1</sub> (Table 1).

**Table 1:** Occurrence of AFM<sub>1</sub> in urine samples of participants according to socio-demographic factors

Socio-demographic factors	Sample tested No.	Positive samples No.	Min- max (pg/ml)	Mean (pg/ml)
<b>Occupational position</b>				
Academic	9	2	0-5.8	0.6
Non- Academic	61	13	0-42	1.7
<b>Age (year)</b>				
20-30	14	2	0-5.8	0.4
31-40	31	9	0-42	2.1
41-50	25	4	0-32.6	1.4
<b>Professional experience (years)</b>				
<5	26	5	0-8.6	0.7
5-10	8	3	0-42	5.5
>10	36	7	0-32.6	1.3
<b>Educational level</b>				
BSc	35	6	0-32.6	1.3
MSc	27	7	0-42	2.1
PhD	8	2	0-5.8	0.7
<b>Sex</b>				
Male	32	7	0-9.2	0.6
Female	38	8	0-42	2.3
<b>Monthly income</b>				
<387\$	23	4	0-9.2	0.6
387-1160 \$	38	9	0-42	2.3
>1160 \$	9	2	0-5.8	0.6
<b>Marital status</b>				
Married	64	12	0-42	1.6
Single	6	3	0-5.8	1.3

**Table 2:** Occurrence of AFM<sub>1</sub> in urine samples of participants according to consumed foods in recent 72 h

Consumed foods in recent 72 h	Sample tested No.	Positive samples No.	Min- Max (pg/ml)	Mean (pg/ml)	P value
<b>Milk</b>					
Yes	64	14	0-42	1.7	0.5
No	6	1	0-0.1	0.1	
<b>Meat</b>					
Yes	62	12	0-32.6	1.1	0.06
No	8	3	0-42	5.6	
<b>Iranian traditional confection</b>					
Yes	16	6	0-42	5	0.01
No	54	9	0-9.2	0.5	
<b>Nuts</b>					
Yes	24	10	0-42	4	0.01
No	45	4	0-3.6	0.1	
<b>Halva Ardeh ( confection made from sesame)</b>					
Yes	5	1	0-1.8	0.3	0.6
No	64	14	0-42	1	
<b>Rice</b>					
Native					
Yes	42	10	0-42	1.9	0.79
No	1	0	0	0	
Imported					
Yes	10	2	0-9.2	1.3	0.72
No	1	0	0	0	
Both					
Yes	16	3	0-8.6	0.9	0.37
No	0	0	0	0	

## Discussion

In this study, AFM<sub>1</sub> was found in 21% of participant's urine samples with the range of excreted AFM<sub>1</sub> at 0–42 pg/ml. Lei et al. (2013) detected no significant difference between the excretion level of AFM<sub>1</sub> and sex that is similar with our investigation in which no significant relationship was observed between all socio-demographic factors and AFM<sub>1</sub> excretion level in urine samples of participants.

Several studies have examined the relationship between food intake and urinary excretion of AFM<sub>1</sub>. Redzwan et al. (2012) found significant relationship between consumption of milk and dairy products and excretion of AFM<sub>1</sub>. In another investigation, Romero et al. (2010) evaluated the presence of AFM<sub>1</sub> in human urine samples from a Brazilian population regarding their corn, peanut, and milk consumption amounts. Their results showed no significant difference between food consumption and excretion of AFM<sub>1</sub> that is in accordance with our findings.

Based on this study, there was not found any significant differences between the excretion level of AFM<sub>1</sub> with the taking some of foods such as milk, meat, rice and Halva Ardeh (a confection made from sesame). But, there was significant association between AFM<sub>1</sub> excretion and consumption of nuts. Several studies have reported that AFB<sub>1</sub> occurs in edible nuts of Iran as high as 77-99% (Cheraghali et al., 2007; Pour et al., 2010; Yazdanpanah et al., 2013) that confirm this finding observed in our survey.

It was observed that the urinary excretion level of AFM<sub>1</sub> in participants who had used traditional confection of Yazd, Iran was significantly higher than those who had not consumed. This matter could be due to considerable AFB<sub>1</sub> contamination rate of major ingredients used in this Iranian traditional product. Wheat flour, walnuts, pistachios and almonds are the main components which are usually used to produce traditional confection of this region of Iran. It seems that due to economic benefit, some Iranian confectioners generally use low quality ingredients which are cheaper than high quality ones. Based on present scientific databases, it has been indicated that low quality nuts and cereal are generally more contaminated to AFB<sub>1</sub> (Afshar et al., 2013; Cheraghali et al., 2007; Feizy et al., 2010; Montaseri et al., 2014; Pour et al., 2010; Sadeghi et al., 2009; Yazdanpanah et al., 2013).

Excretion of AFM<sub>1</sub> in this study was lower than in countries such as Brazil, Egypt (Iha et al., 2011; Polychronaki et al., 2008). It should be noted that minimum education level of all participants in this survey was BSc. It can be assumed that individuals with higher education level may be more awarded and knowledgeable

about AF contamination and its implication for human health, and this is probably the main reason for the low excretion of AFM<sub>1</sub> the participants of this survey.

Previous studies have shown that probiotic bacteria could reduce the detrimental effects of AF (Byun and Yoon, 2003; Hernandez Mendoza et al., 2009; Montaseri et al., 2014). Therefore, the consumption of probiotic yogurt should be increased to minimize the adverse health effects of the toxin. Also, some researchers have indicated that using clay, as a good adsorbent for AF in intestinal, could effectively reduce human exposure to AF (Phillips et al., 2008). However, it seems that comprehensive training of the people is the most practical and useful method for this proposes.

## Conclusion

Excretion level of AFM<sub>1</sub>, as a biomarker of AFB<sub>1</sub> exposure, in urine samples of Iranian population was evaluated. Our findings indicate that exposure to the AF in staffs of Faculty of Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran was relatively notable. Considering severe adverse health effects of AF, the state of a comprehensive plan for training of the people to avoid eating high contaminated food, is one of the most practical and useful methods for reducing AF health risks.

## Conflicts of interest

The authors declare that they had no conflict of interest.

## Acknowledgement

This work was funded by Shahid Sadoughi University of Medical Sciences, Yazd, Iran. We thank Mr. Iman Moini-Nasab and Mr. M. Mohammadzadeh for their kindly help during sampling and data analysis.

## References

- Afshar P., Shokrzadeh M., Kalthori S., Babbai Z. (2013). Occurrence of ochratoxin A and aflatoxin M<sub>1</sub> in human breast milk in Sari, Iran. *Food Control*. 31: 525-529.
- Asadi M., Beheshti H.R., Feizy J. (2011). A survey of aflatoxins in sesame in Iran. *Mycotoxin research*. 27: 259-263.
- Atanda O., Akpan I., Oluwafemi F. (2007). The potential of some spice essential oils in the control of *A. parasiticus* CFR 223 and aflatoxin production. *Food Control*. 18: 601-607.
- Battacone G., Nudda A., Palomba M., Pascale M. (2005). Transfer of aflatoxin B<sub>1</sub> from feed to milk and from milk to curd and whey in dairy sheep fed artificially contaminated concentrates. *Journal of Dairy Science*. 88: 3063-3069.
- Byun J.R., Yoon Y.H. (2003). Binding of aflatoxin G<sub>1</sub>, G<sub>2</sub> and B<sub>2</sub> by probiotic *Lactobacillus* spp. *Asian-Australasian Journal of Animal Science*. 16: 1686-1689.
- Cheraghali A., Yazdanpanah H., Abouhossain G., Doraki N. (2007). Incidence of aflatoxins in Iran pistachio nuts. *Food and Chemical Toxicology*. 45: 812-816.

- Egal S., Hounsa A., Gong Y., Turner P., Wild C. (2005). Dietary exposure to aflatoxin from maize and groundnut in young children from Benin and Togo, West Africa. *International Journal of Food Microbiology*. 104: 215-224.
- Ehrlich K.C., Kobbeman K., Montalbano B.G. Cotty P.J. (2007). Aflatoxin-producing *Aspergillus* species from Thailand. *International Journal of Food Microbiology*. 114: 153-159.
- Feizy J., Beheshti H., Fahimi N.K., Janati S.F. (2010). Survey of aflatoxins in rice from Iran using immunoaffinity column clean-up and HPLC with fluorescence detection. *Food Additives and Contaminants*. 3: 263-267.
- Groopman J.D., Kensler T.W. (2005). Role of metabolism and viruses in aflatoxin-induced liver cancer. *Toxicology and Applied Pharmacology*. 206: 131-137.
- Hernandez Mendoza A., Guzman De Pena D., Garcia H.S. (2009). Key role of teichoic acids on aflatoxin B<sub>1</sub> binding by probiotic bacteria. *Journal of Applied Microbiology*. 107: 395-403.
- Iha M.H., Barbosa C.B., Okada I.A. (2011). Occurrence of aflatoxin M<sub>1</sub> in dairy products in Brazil. *Food Control*. 22: 1971-1974.
- Khodadadi M., Khosravi R., Allahresani A., Khksar Y., Rafati L. (2014). Occurrence of aflatoxin M<sub>1</sub> in pasteurized and traditional cheese marketed in southern Khorasan, Iran. *Journal of Food Quality and Hazards Control*. 1: 77-80.
- Lei Y., Fang L., Akash M.S.H., Rehman K., Liu Z., Shi W., Chen S. (2013). Estimation of urinary concentration of aflatoxin M<sub>1</sub> in Chinese pregnant women. *Journal of Food Science*. 78: 1835-1838.
- Leong Y.H., Latiff A.A., Ahmad N.I., Rosman A. (2012). Exposure measurement of aflatoxins and aflatoxin metabolites in human body fluids. A short review. *Mycotoxin Research*. 28: 79-87.
- Lewis L., Onsongo M., Njapau H., Schurz H., Lubner G., Kieszak S. (2005). Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in eastern and central Kenya. *Environmental Health Perspectives*. 113: 1763-1767.
- Mahdavi R., Nikniaz L., Arefhosseini S., Jabbari M.V. (2010). Determination of aflatoxin M<sub>1</sub> in breast milk samples in Tabriz-Iran. *Maternal and Child Health Journal*. 14: 141-145.
- McKean C., Tang L., Tang M., Billam M., Wang Z. (2006). Comparative acute and combinative toxicity of aflatoxin B<sub>1</sub> and fumonisin B<sub>1</sub> in animals and human cells. *Food and Chemical Toxicology*. 44: 868-876.
- Montaseri H., Arjmandtalab S., Dehghanzadeh G., Karami S., Razmjoo M., Sayadi M., Oryan A. (2014). Effect of production and storage of probiotic yogurt on aflatoxin M<sub>1</sub> residue. *Journal of Food Quality and Hazards Control*. 1: 7-14.
- Partanen H.A., EL-Nezami H.S., Leppanen J.M. (2009). Aflatoxin B<sub>1</sub> transfer and metabolism in human placenta. *Toxicological Sciences*. 113: 216-225.
- Phillips T., Afriyie-Gyawu E., Williams J., Huebner H., Ankrah N.A., Ofori-Adjei D., Jolly P. (2008). Reducing human exposure to aflatoxin through the use of clay: a review. *Food Additives and Contaminants*. 25: 134-145.
- Polychronaki N., Wild C.P., Mykkänen H., Amra H., Abdel-Wahhab M., Sylla A., Diallo M. (2008). Urinary biomarkers of aflatoxin exposure in young children from Egypt and Guinea. *Food and Chemical Toxicology*. 46: 519-526.
- Pour R.S., Rasti M., Zighamian H., Garmakhani A.D. (2010). Occurrence of aflatoxins in pistachio nuts in Esfahan Province of Iran. *Journal of Food Safety*. 30: 330-340.
- Rahimi E., Bonyadian M., Rafei M., Kazemian H. (2010). Occurrence of aflatoxin M<sub>1</sub> in raw milk of five dairy species in Ahvaz, Iran. *Food and Chemical Toxicology*. 48: 129-131.
- Redzwan S.M., Rosita J., Mohd Sokhini A.M., Nurul Aqilah A.R. (2012). Association between aflatoxin M<sub>1</sub> excreted in human urine samples with the consumption of milk and dairy products. *Bulletin of Environmental Contamination and Toxicology*. 89: 1115-1119.
- Romero A., Ferreira T., Dias C., Calori-Domingues M.A. (2010). Occurrence of AFM<sub>1</sub> in urine samples of a Brazilian population and association with food consumption. *Food Control*. 21: 554-558.
- Sabran M.R., Jamaluddin R., Abdulmutalib M.S. (2012). Screening of aflatoxin M<sub>1</sub>, a metabolite of aflatoxin B<sub>1</sub> in human urine samples in Malaysia: a preliminary study. *Food Control*. 28: 55-58.
- Sadeghi N., Ovesi M.R., Jannat B., Hajjimaahmoodi M., Bonyani H., Jannat F. (2009). Incidence of aflatoxin M<sub>1</sub> in human breast milk in Tehran, Iran. *Food Control*. 20: 75-78.
- Sherif S.O., Salama E.E., Abdel M.A. (2009). Mycotoxins and child health: the need for health risk assessment. *International Journal of Hygiene and Environmental Health*. 212: 347-368.
- Shuaib F., Ehiri J., Abdullahi A., Williams J.H., Jolly P.E. (2010). Reproductive health effects of aflatoxins: a review of the literature. *Reproductive Toxicology*. 29: 262-270.
- Smith L.E., Stoltzfus R.J., Prendergast A. (2012). Food chain mycotoxin exposure, gut health, and impaired growth: a conceptual framework. *Advances in Nutrition*. 3: 526-531.
- Wu H.C., Wang Q., Yang H.I., Ahsan H., Tsal W.Y., Wang L.Y., Chen S.Y., Chen C.J., Santella R.M. (2009). Aflatoxin B<sub>1</sub> exposure, hepatitis B virus infection, and hepatocellular carcinoma in Taiwan. *Cancer Epidemiology Biomarkers and Prevention*. 18: 846-853.
- Yazdanpanah H., Zarghi A., Shafaati A.R., Foroutan S.M., Aboul-Fathi F., Khoddam A., Nazari F. (2013). Analysis of aflatoxin B<sub>1</sub> in Iranian foods using HPLC and a monolithic column and estimation of its dietary intake. *Iranian Journal of Pharmaceutical Research*. 12: 83-89.
- Zaghini A., Martelli G., Roncada P., Simoli M., Rizzi L. (2005). Mannan oligosaccharides and aflatoxin B<sub>1</sub> in feed for laying hens: effects on egg quality, aflatoxins B<sub>1</sub> and M<sub>1</sub> residues in eggs, and aflatoxin B<sub>1</sub> levels in liver. *Poultry Science*. 84: 825-832.