



Editorial

Special Attention is Needed for Reduce Antibiotic Residue Risk in the White Meat Produced in Algeria

A. Berghiche

Laboratory of Life Sciences and Techniques, Institute of Agronomic and Veterinary Sciences, Mohamed Cherif Messaadia University, Souk Ahras, Algeria

* Corresponding author. ✉ amine_berghiche@yahoo.com
ORCID ID: <https://orcid.org/0000-0001-6355-3591>

The therapeutic use of antibiotics in human and veterinary medicine constituted an effective weapon for a long time counters many pathogenic germs. However, abusive use of certain antibiotics in preventive treatment or as a food additive led to possibilities of development of some health problems such as the allergic reactions, cancer, and even increasing the populations of antibiotics resistant microbes (Berghiche et al., 2017; Nisha, 2008; Zhang et al., 2003). Abuse of antibiotics is a common practice in the field of poultry farming in Algeria, while farmers are usually unaware of the risks associated with the antibiotics residue and do not often attention to withdrawal period for each antibiotic (Berghiche et al., 2018; Goossens, 2014). If the antibiotics used by veterinarians and poultry farmers exceed, the limits consumers' health may be endangered. There is a great lack of health monitoring about abuse of antibiotics in Algerian poultry farms. So, because of health side effects of antibiotic residue in white meat produced in Algeria, implementation of effective programs is too vital by national

authorities to reduce this hygienic risk.

References

- Berghiche A., Khenenou T., Bouzebda-Afri F., Lamraoui R., Labiad I. (2017). Detection of the antibiotic residues in broiler chickens by microbiological screening test in Algeria. *Global Veterinaria*. 19: 504-508. [DOI: 10.5829/idosi.gv.2017.504.508]
- Berghiche A., Khenenou T., Kouzi A., Labiad I. (2018). An investigation on the predominant diseases, its diagnosis, and commonly used drugs in the poultry farms in the North-Eastern regions of Algeria. *Veterinary World*. 11: 986-989. [DOI: 10.14202/vetworld.2018.986-989]
- Goossens H. (2014). European strategies to control antibiotic resistance and use. *Annals of Clinical Microbiology*. 17: 1-8. [DOI: 10.5145/ACM.2014.17.1.1]
- Nisha A.R. (2008). Antibiotic residues- a global health hazard. *Veterinary World*. 1: 375-377.
- Zhang Q., Lin J., Pereira S. (2003). Fluoroquinolone-resistant *Campylobacter* in animal reservoirs: dynamics of development, resistance mechanisms and ecological fitness. *Animal Health Research Reviews*. 4: 63-72. [DOI:10.1079/AHRR.200356]

© 2019, Shahid Sadoughi University of Medical Sciences. This is an open access article under the Creative Commons Attribution 4.0 International License.

To cite: Berghiche A. (2019). Special attention is needed for reduce antibiotic residue risk in the white meat produced in Algeria. *Journal of Food Quality and Hazards Control*. 6: 44.