



Dominant Lactic Acid Bacteria in Naturally Fermented Milks from Messinese Goat's Breed

M. Palmeri¹, I. Mancuso¹, P. Barbaccia², F. Cirlincione², M.L. Scatassa^{1*}

1. Istituto Zooprofilattico Sperimentale della Sicilia "Adelmo Mirri", Via G. Marinuzzi 3, 90129 Palermo, Italy

2. Dipartimento Scienze Agrarie, Alimentari e Forestali, Università di Palermo, Viale delle Scienze 4, 90128 Palermo, Italy

HIGHLIGHTS

- Levels of coccus- and rod-shaped LAB were in the range of 1.78-7.10 log and 1.00-7.09 log colony forming unit/ml, respectively.
- Among 12 identified strains, the most numerous one was *Enterococcus hirae* (n=4) followed by *E. faecium* (n=3).
- Risk assessment of pathogenic enterococci species is needed for the consumers of traditional fermented dairy products.

Article type

Original article

Keywords

Milk
Goats
Lactobacillales
Cultured Dairy Products
Food Microbiology

Article history

Received: 18 Jan 2019

Revised: 27 Mar 2019

Accepted: 11 Apr 2019

Acronyms and abbreviations

LAB=Lactic Acid Bacteria
RAPD=Random Amplification of Polymorphic DNA
PCR=Polymerase Chain Reaction
CFU=Colony Forming Unit
MRS=de Man-Rogosa-Sharpe
M17=Medium 17 agar
FGM=Fermented Goat's Milk

ABSTRACT

Background: Lactic Acid Bacteria (LAB) are an important group of microorganisms responsible for the fermentation dairy products. This study was done to identify the dominant lactic acid bacteria in naturally fermented milks from Messinese goat's breed.

Methods: Eighteen individual raw milk samples collected from Messinese goat's breed were acidified at pH 5.20 and left to spontaneously ferment at 37 °C for 4 days. All samples were analyzed for rod- and coccus-shaped LAB. Also, all presumptive LAB were isolated and differentiated according to their phenotypic properties and genetic polymorphisms and then identified by sequencing the 16S rRNA gene. Data were statistically analyzed using SAS 9.2 software.

Results: Levels of coccus- and rod-shaped LAB were in the range of 1.78-7.10 log and 1.00-7.09 log colony forming unit/ml, respectively. The microbiological counts on the two different growth media were significantly ($p<0.05$) different among the samples. Among 12 identified strains, the most numerous one was *Enterococcus hirae* (n=4), followed by *E. faecium* (n=3), while the other species (*E. durans*, *E. faecalis*, *E. lactis*, *Lactococcus lactis*, and *Leuconostoc lactis*) included one strain each.

Conclusion: The major group identified in this study was mainly represented by members of *Enterococcus* genus. Although *Enterococcus* spp. are related to the typicality of some traditional fermented dairy products, this study highlights the need for risk assessment of pathogenic enterococci species for the consumers.

© 2019, Shahid Sadoughi University of Medical Sciences. This is an open access article under the Creative Commons Attribution 4.0 International License.

Introduction

Lactic Acid Bacteria (LAB) are an important group of microorganisms responsible for the fermentation of a large variety of foods, including those derived from dairy products (Gaglio et al., 2014a; Gaglio et al., 2019a). LAB

are able to improve the final quality of dairy productions and in particular their flavour, texture, and nutritional value (Gaglio et al., 2016a; Guarrasi et al., 2017; Leroy and De Vuyst, 2004). Furthermore, they have the ability

* Corresponding author. ✉ luisa.scatassa@izssicilia.it
ORCID ID: <http://orcid.org/0000-0001-5132-8768>

To cite: Palmeri M., Mancuso I., Barbaccia P., Cirlincione F., Scatassa M.L. (2019). Dominant lactic acid bacteria in naturally fermented milks from Messinese goat's breed. *Journal of Food Quality and Hazards Control*. 6: 66-72.

to determine the rapid acidification of the raw materials and inhibit the growth of spoilage and pathogenic microorganisms (Guarcello et al., 2016; Macaluso et al., 2016; Scatassa et al., 2017; Settanni et al., 2013).

The use of starter LAB to ferment milk with particular nutritional properties (e.g. goat's milk), represents one of the technological strategy to develop a large variety of new dairy functional foods and beverages (Minervini et al., 2009). Goat's milk is well recognized as "the king of milk" for its high digestibility and nutritional value as well as for the lower allergenic compound content than cow's milk (Ribeiro and Ribeiro, 2010). Furthermore, goat's milk is one of the matrices most closed to human milk and plays an important role in healthy diet of children and elderly people (Haenlein, 2004).

Although several works are available on the characterization of LAB from goat's cheeses and fermented milks in literature (Meng et al., 2018; Minervini et al., 2009), only a few works focused on the characteristics of the raw milks before processing. In order to develop *ad hoc* starter cultures for a given product, the characterization of the populations associated with raw materials is of relevance. The particular flavor and typical organoleptic properties of products processed from raw milk are related not only to the race and nutrition of animals, but also to the natural microbiota responsible for fermentation (Franciosi et al., 2008). With this in mind, the present work was carried out to identify the autochthonous LAB composition of naturally Fermented Goat's Milks (FGM) in order to characterize the dominant microbial populations.

Materials and methods

Sample collection, acidification, and incubation of goat's milk

Eighteen individual milk samples of Messinese goat's breed were collected from a farm located in Palermo province, Italy. Just after sampling, all samples were placed into a portable fridge and transferred to the Laboratory of Centro Latte e Lotta alle Mastiti (Istituto Zooprofilattico Sperimentale della Sicilia "Adelmo Mirri", Palermo, Italy) where they were immediately subjected to the acidification procedure; in order to allow exclusively the growth of LAB, each milk sample was added with 5 M lactic acid (BDH Prolabo Chemicals, Singapore) until reaching the pH 5.20 evaluated electrometrically by pH meter HI3220-02 (Hanna Instruments, Woonsocket, RI, USA). All samples were incubated at 37 °C for 4 days. The milk samples from each goat were collected in duplicate at two-week intervals.

Microbiological analyses, LAB isolation, and phenotypic grouping

All 18 FGM samples (FMG1-FMG18) were serially diluted in Ringer's solution (Sigma-Aldrich, Milan, Italy). Cell suspensions were then subjected to plate count for the enumeration of rod- and coccus-shaped LAB after pouring on de Man-Rogosa-Sharpe (MRS) agar acidified with 5 M lactic acid to pH 5.4 incubated anaerobically at 37 °C for 48 h and on Medium 17 (M17) agar incubated aerobically at 37 °C for 48 h, respectively (Aureli et al., 2008). Incubation of rod-shaped LAB was occurred in anaerobiosis using the AnaeroGen AN25 (Oxoid, Milan, Italy) in jars closed hermetically. Both MRS and M17 were purchased from Oxoid®, UK.

After growth, colonies of various shapes of Gram-positive and catalase negative bacteria were randomly picked from count plates considering all different colours, edges, and elevations, then transferred into the corresponding broth media. All different morphologies were considered in order to evaluate total LAB diversity. The isolates were purified by successive subculturing and stored in MRS or M17 broth media containing 20% glycerol (v/v) at -80 °C until further analysis.

All presumptive LAB isolates from FGM were subjected to a phenotypic characterization on the basis of cell morphology, cell disposition, growth at 15 and 45 °C, resistance at 60 °C for 30 min, NH₃ production from arginine, aesculin hydrolysis, acid production from the carbohydrates (arabinose, ribose, xylose, fructose, galactose, lactose and sucrose), and CO₂ production from glucose (Di Grigoli et al., 2015; Gaglio et al., 2014b). The coccus-shaped isolates were further grouped by their ability to grow at pH 9.2 and in the presence of NaCl (6.5 g/l) to separate enterococci from other dairy cocci.

Genotypic differentiation and identification of bacteria

DNAs from LAB cultures were extracted using the InstaGene Matrix kit (Bio-Rad, Hercules, CA, USA) following the manufacturer's instructions. Cells were harvested after overnight growth in MRS or M17 broths at 37 °C and washed in distilled H₂O; then the crude cell extracts were used as templates for Polymerase Chain Reactions (PCRs).

The differentiation of the isolates at strain level was performed by Random Amplification of Polymorphic DNA (RAPD)-PCR analysis in a 30 µl reaction volume with the primers M13, AB111, and AB106 used singly as reported by Gaglio et al. (2017). PCR mixture included 62.5 ng of target DNA, 2.5 µl of PCR buffer (Fermentas, MMedical, Milan, Italy), 2.5 mM of MgCl₂, 250 µM of each dNTP (Life Technologies Monza, Italy), 0.2 µM of each primer, 2.5 U of Taq DNA polymerase

(Thermo Fisher Scientific, Monza, Italy), and Milli-Q water to reach the final reaction volume. The PCR program applied for all primers comprised 40 cycles of denaturation for 2 min at 94 °C, annealing for 20 s at 40 °C, and extension for 2 min at 72 °C; the cycles were preceded by denaturation at 94 °C for 2 min and followed by extension at 72 °C for 5 min. The amplifications were performed using a Thermal cycler (Bioer, Hangzhou, China) and the amplified products were separated by electrophoresis, visualized and acquired by Gel Doc™ XR+ and ChemiDoc™ XRS+ Imaging Systems (Bio Rad, Hercules, CA, USA). The analysis of the RAPD patterns was performed with the Gelcompar II software, version 6.5 (Applied-Maths, Sint-Martens-Latem, Belgium).

All LAB showing different RAPD-PCR profiles were genetically identified at species level by 16S rRNA gene sequencing as described by Weisburg et al. (1991). PCR mixture (50 µl total volume) included 62.5 ng of target DNA, 1×Taq DNA polymerase buffer with 2 mM MgCl₂ (Thermo Fisher Scientific, Monza, Italy), 250 µM of each dNTP (Life Technologies, Italy), 0.2 µM of each primer, 2.5 U of Taq DNA polymerase (Thermo Fisher Scientific, Monza, Italy), and Milli-Q water to reach the final reaction volume. PCR program comprised an initial template denaturation step for 3 min at 95 °C followed by 30 cycles of denaturation for 1 min at 94 °C, annealing for 45 s at 54 °C, and extension for 2 min at 72 °C. The final elongation step was for 7 min at 72 °C. The PCR products were visualized as reported above and the amplicons corresponding approximately 1600 bp were purified using the Illustra GFX PCR DNA and Gel Band Purification Kit (GE Healthcare Bio-Sciences, Pittsburgh, PA, USA). Sequencing analyses were performed in an ABI PRISM 3500 Genetic Analyzer (Applied Biosystems, Carlsbad, CA, USA) and compared using a BLAST search in the GenBank/ EMBL/DDBJ database (<http://www.ncbi.nlm.nih.gov>).

Statistical analyses

Statistical analyses of microbiological data were subjected to one-way Analysis of Variance (ANOVA) using SAS 9.2 software (Statistical Analysis System Institute Inc., Cary, NC, USA). Pair comparison between means were determined by the post-hoc Duncan test at $p < 0.05$. Differences between the samples were analyzed using a Generalised Linear Model (GLM) procedure.

Results

Figure 1 shows the viable counts of the rod- and coccus-shaped LAB harboured on FGM samples. Levels of these microbial groups for each sample were almost comparable. The microbiological counts on the two

different growth media were significantly ($p < 0.05$) different among the samples. In fact, the levels of coccus-shaped LAB were in the range of 1.78-7.10 log Colony Forming Unit (CFU)/ml while rod-shaped LAB were in the range of 1.00-7.09 log CFU/ml. Samples FGM1, FGM4, FGM9, FGM14, FGM16, FGM17, and FGM18 displayed the highest bacterial density of rod- and coccus-shaped LAB; while the samples FM8 and FM11 showed the lowest levels of these bacterial groups.

A total of 247 colonies were collected from the 18 samples of FGM. After purification and microscopic inspection, all the cultures showed a coccus-shaped morphology. One hundred eighty seven cultures were considered presumptive LAB, being Gram-positive and catalase negative. According to the combination of the phenotypic properties evaluated the 187 presumptive LAB cultures, were separated into five groups (Table 1). The most numerous groups were groups II and III, composed of more than 50 isolates each. LAB cultures included between the groups I to IV showed an obligate homofermentative metabolism while group V showed a heterofermentative metabolism.

Approximately 30% of the isolates from each phenotypic group, forming a total of 56 isolates, were subjected to RAPD-PCR analysis for strain typing. The genotyping differentiation indicated that the cultivable bacterial community associated to FGM in the present study was composed of 12 distinct strains (Figure 2). The dendrogram clearly showed that the strains belonging to the phenotypic groups I to III clustered closely after RAPD-PCR analysis.

Twelve strains were identified by sequencing of the 16S rRNA gene. The sequence comparison within BLAST database identified seven major dominating species. The species with the highest number of strains was *Enterococcus hirae* (n=4) followed by *E. faecium* (n=3), while the other species (*E. durans*, *E. faecalis*, *E. lactis*, *Lactococcus lactis*, and *Leuconostoc lactis*) included one strain each.

Discussion

Some previous works about LAB in raw ewe's and cow's milk showed potential of LAB to drive the fermentation/ripening processes of different dairy products (Gaglio et al., 2014b; Guarcello et al., 2016; Turchi et al., 2011); however, few information are available on goat's LAB from Italian milk products. In the present study, all goat milk samples were dominated by rod- and coccus-shaped LAB, but only a few samples showed levels comparable with those of typical milk based fermented products, such as yogurt (Rezac et al., 2018) with the levels at around 7 log CFU/ml. The presumptive LAB

Table 1: Phenotypic grouping of the LAB isolated from fermented goat's milks

Characteristics	Clusters				
	I (n=13)	II (n=62)	III (n=81)	IV (n=19)	V (n=12)
Morphology	C*	C	C	C	C
Cell disposition	SC**	SC	SC	SC	SC
Growth:					
15 °C	+	+	+	+	+
45 °C	+	+	+	-	+
pH 9.2	+	+	+	+	-
6.5% NaCl	+	+	+	-	+
Resistance to 60 °C					
	-	-	+	+	-
Hydrolysis of:					
Arginine	+	+	+	+	-
Aesculin	+	+	+	+	-
Acid production from:					
Arabinose	+	+	+	-	+
Ribose	+	+	+	+	+
Xylose	+	+	+	-	+
Fructose	+	+	+	+	+
Galactose	+	+	+	+	+
Lactose	+	+	+	+	+
Sucrose	+	-	+	+	+
Glycerol	+	+	+	+	+
CO ₂ from glucose	-	-	-	-	+

C: Coccus; SC: Short Chain




Figure 1: Microbiological concentrations (CFU; Colony Forming Unit/ml) of fermented goat's milk samples. MRS: de Man-Rogosa-Sharpe agar for detection of rod-shaped LAB; M17: Medium 17 agar for detection of coccus-shaped LAB; FGM: Fermented Goat's Milk. Different superscript letters indicate significant differences on microbial loads were performed for each sample according to Duncan test between MRS and M17 media for $p < 0.05$




Figure 2: Dendrogram obtained with combined RAPD-PCR patterns of the LAB strains isolated from fermented goat's milks. Scale bar indicate the percentage of similarity. *E.*: *Enterococcus*; *Lc.*: *Lactococcus*; *Ln.*: *Leuconostoc*

isolated from the FGM samples represented five phenotypic groups of cocci, even though on MRS should preferably develop rod-shaped species. However, despite the high count detected in MRS, our results confirmed that LAB cocci are able to develop on this media commonly used for mesophilic as well as thermophilic rods LAB, maybe due to the lower pH of this media (Settanni et al., 2012).

RAPD-PCR, a technique commonly used to differentiate microorganisms indicated the presence of 12 different strains in our goat milk samples. This biodiversity is quite low compared to that of traditional cheese evidenced by Franciosi et al. (2008), but the explanation for this controversy could be due to the fact that only one farm was analyzed. In a single farm, even if several animals are object of investigation, the environmental factors and the feed sources are the same and this might have reduced biodiversity found among milk samples. The genetic identification resulted in the following LAB species: *E. durans*, *E. faecalis*, *E. faecium*, *E. hirae*, *E. lactis*, *Lc. Lactis*, and *Ln. lactis*. All these species are commonly associated with raw milk and cheeses (Franciosi et al., 2011; Gaglio et al., 2019b) and the equipment commonly used in traditional cheese productions (Cruciata et al., 2018; Scatassa et al., 2015). Among them, *Lc. lactis* and *Ln. lactis* are typical mesophilic starter cultures for dairy use (Settanni and Moschetti,

2010), while the other identified species are commonly shown to be part of the non-starter LAB population implicated in the maturation of several cheeses (Gatti et al., 2014; Settanni and Moschetti, 2014).

In the current research, the high number of strains detected as members of the *Enterococcus* spp. was surprising especially those belonging to the species *E. hirae* and *E. faecium*. Although in other studies, enterococci were not found at consistent levels in FGM, their presence at 10^6 – 10^8 CFU/g was reported by some researchers who analyzed the traditional ovine and caprine cheeses produced in Greece (Nikolaou et al., 2002; Prodromou et al., 2001; Psoni et al., 2003). It is well known that enterococci are an integral part of the microbial population of different typologies of traditional dairy productions (Nikolaou et al., 2002; Psoni et al., 2003; Suzzi et al., 2000) and due to their proteolytic and lipolytic activities, they are involved in the development of typicality organoleptic characteristics (Giraffa, 2002). On the other hand, it should be highlighted that in the last years, enterococci are considered as emerging pathogens for humans may endangering public health. So, inclusion of enterococci as starter cultures for cheese production needs to be validated by the absence of pathogenic traits, such as antibiotic resistance, virulence and haemolytic activity (Gaglio et al., 2016b; Russo et al., 2018; Settanni et al., 2014).

Conclusion

This work showed the importance of selecting acidifying LAB from raw materials for production of fermented dairy products; since they might ensure the ability of persist during fermentation. The major group identified in this study was mainly represented by members of *Enterococcus* genus. Although enterococci sometimes are related to the typicality of some traditional fermented dairy products, this study highlights the need for risk assessment of pathogenic enterococci species for the consumers.

Author contributions

M.L.S. designed the project of study; P.B., F.C., and M.P. conducted the experiments; M.P. and I.M. analyzed the data; M.L.S. and I.M. wrote the manuscript. All authors revised and approved the final manuscript.

Conflicts of interest

There was no conflict of interest in this study.

Acknowledgements

This work was financed by the Italian Ministry of Health Research Project RC 06/16 Valutazione del rischio associato alla presenza di enterococchi antibiotico-resistenti nei prodotti lattiero-caseari e sviluppo di metodologie che consentano la preservazione delle produzioni (CUP H76J17000480001). The authors are grateful to Dr. Rossella Calascibetta for their precious collaboration during samples collection.

References

- Aureli P., Fiore A., Scalfaro C., Franciosa G. (2008). Microbiological and molecular methods for analysis of probiotic based food supplements for human consumption. *Rapporti Istisan*.
- Cruciata M., Gaglio R., Scatassa M.L., Sala G., Cardamone C., Palmeri M., Moschetti G., La Mantia T., Settanni L. (2018). Formation and characterization of early bacterial biofilms on different wood typologies applied in dairy production. *Applied and Environmental Microbiology*. 84: e02107-17. [DOI: 10.1128/AEM.02107-17]
- Di Grigoli A., Francesca N., Gaglio R., Guarrasi V., Moschetti M., Scatassa M.L., Settanni L., Bonanno A. (2015). The influence of the wooden equipment employed for cheese manufacture on the characteristics of a traditional stretched cheese during ripening. *Food Microbiology*. 46: 81-91. [DOI: 10.1016/j.fm.2014.07.008]
- Franciosi E., Settanni L., Carlin S., Cavazza A., Poznanski E. (2008). A factory-scale application of secondary adjunct cultures selected from lactic acid bacteria during Puzzone di Moena cheese ripening. *Journal of Dairy Science*. 91: 2981-2991. [DOI: 10.3168/jds.2007-0764]
- Franciosi E., Settanni L., Cologna N., Cavazza A., Poznanski E. (2011). Microbial analysis of raw cows' milk used for cheese-making: influence of storage treatments on microbial composition and other technological traits. *World Journal of Microbiology and Biotechnology*. 27: 171-180. [DOI: 10.1007/s11274-010-0443-2]
- Gaglio R., Couto N., Marques C., Lopes M.D.F.S., Moschetti G., Pomba C., Settanni L. (2016b). Evaluation of antimicrobial resistance and virulence of enterococci from equipment surfaces, raw materials, and traditional cheeses. *International Journal of Food Microbiology*. 236: 107-114. [DOI: 10.1016/j.ijfoodmicro.2016.07.020]
- Gaglio R., Cruciata M., Di Gerlando R., Scatassa M.L., Cardamone C., Mancuso I., Sardina M.T., Moschetti G., Portolano B., Settanni L. (2016a). Microbial activation of wooden vats used for traditional cheese production and evolution of neofomed biofilms. *Applied and Environmental Microbiology*. 82: 585-595. [DOI: 10.1128/AEM.02868-15]
- Gaglio R., Cruciata M., Scatassa M.L., Tolone M., Mancuso I., Cardamone C., Corona O., Todaro M., Settanni L. (2019b). Influence of the early bacterial biofilms developed on vats made with seven wood types on PDO Vastedda della valle del Belice cheese characteristics. *International Journal of Food Microbiology*. 291: 91-103. [DOI: 10.1016/j.ijfoodmicro.2018.11.017]
- Gaglio R., Francesca N., Di Gerlando R., Cruciata M., Guarcello R., Portolano B., Moschetti G., Settanni L. (2014b). Identification, typing, and investigation of the dairy characteristics of lactic acid bacteria isolated from "Vastedda della valle del Belice" cheeses. *Dairy Science and Technology*. 94: 157-180. [DOI: 10.1007/s13594-013-0150-5]
- Gaglio R., Francesca N., Di Gerlando R., Mahony J., De Martino S., Stucchi C., Moschetti G., Settanni L. (2017). Enteric bacteria of food ice and their survival in alcoholic beverages and soft drinks. *Food Microbiology*. 67: 17-22. [DOI: 10.1016/j.fm.2017.04.020]
- Gaglio R., Gentile C., Bonanno A., Vintaloro L., Perrone A., Mazza F., Barbaccia P., Settanni L., Di Grigoli A. (2019a). Effect of saffron addition on the microbiological, physicochemical, antioxidant and sensory characteristics of yoghurt. *International Journal of Dairy Technology*. 72: 208-217. [DOI: 10.1111/1471-0307.12569]
- Gaglio R., Scatassa M.L., Cruciata M., Miraglia V., Corona O., Di Gerlando R., Portolano B., Moschetti G., Settanni L. (2014a). *In vivo* application and dynamics of lactic acid bacteria for the four-season production of Vastedda-like cheese. *International Journal of Food Microbiology*. 177: 37-48. [DOI: 10.1016/j.ijfoodmicro.2014.02.007]
- Gatti M., Bottari B., Lazzi C., Neviani E., Mucchetti G. (2014). Invited review: microbial evolution in raw-milk, long-ripened cheeses produced using undefined natural whey starters. *Journal of Dairy Science*. 97: 573-591. [DOI: 10.3168/jds.2013-7187]
- Giraffa G. (2002). Enterococci from foods. *FEMS Microbiology Reviews*. 26: 163-171. [DOI: 10.1111/j.1574-6976.2002.tb00608.x]
- Guarcello R., Carpino S., Gaglio R., Pino A., Rapisarda T., Caggia C., Marino G., Randazzo C.L., Settanni L., Todaro M. (2016). A large factory-scale application of selected autochthonous lactic acid bacteria for PDO Pecorino Siciliano cheese production. *Food Microbiology*. 59: 66-75. [DOI: 10.1016/j.fm.2016.05.011]
- Guarrasi V., Sannino C., Moschetti M., Bonanno A., Di Grigoli A., Settanni L. (2017). The individual contribution of starter and non-starter lactic acid bacteria to the volatile organic compound composition of Caciocavallo Palermitano cheese. *International Journal of Food Microbiology*. 259: 35-42. [DOI: 10.1016/j.ijfoodmicro.2017.07.022]
- Haenlein G.F.W. (2004). Goat milk in human nutrition. *Small Ruminant Research*. 51: 155-163. [DOI: 10.1016/j.smallrumres.2003.08.010]
- Leroy F., De Vuyst L. (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology*. 15: 67-78. [DOI: 10.1016/j.tifs.2003.09.004]

- Macaluso G., Fiorenza G., Gaglio R., Mancuso I., Scatassa M.L. (2016). *In vitro* evaluation of bacteriocin-like inhibitory substances produced by lactic acid bacteria isolated during traditional Sicilian cheese making. *Italian Journal of Food Safety*. 5: 5503. [DOI: 10.4081/ijfs.2016.5503]
- Meng Z., Zhang L., Xin L., Lin K., Yi H., Han X. (2018). Technological characterization of *Lactobacillus* in semihard artisanal goat cheeses from different Mediterranean areas for potential use as nonstarter lactic acid bacteria. *Journal of Dairy Science*. 101: 2887-2896. [DOI: 10.3168/jds.2017-14003]
- Minervini F., Bilancia M.T., Siragusa S., Gobbetti M., Caponio F. (2009). Fermented goats' milk produced with selected multiple starters as a potentially functional food. *Food Microbiology*. 26: 559-564. [DOI: 10.1016/j.fm.2009.03.008]
- Nikolaou E., Tzanetakis N., Litopoulou-Tzanetaki E., Robinson R.K. (2002). Changes in the microbiological and chemical characteristics of an artisanal, low-fat cheese made from raw ovine milk during ripening. *International Journal of Dairy Technology*. 55: 12-17. [DOI: 10.1046/j.1471-0307.2002.00032.x]
- Prodromou K., Thasitou P., Haritonidou E., Tzanetakis N., Litopoulou-Tzanetaki E. (2001). Microbiology of "Orinotyri", a ewe's milk cheese from the Greek mountains. *Food Microbiology*. 18: 319-328. [DOI: 10.1006/fmic.2001.0403]
- Psoni L., Tzanetakis N., Litopoulou-Tzanetaki E. (2003). Microbiological characteristics of Batzos, a traditional Greek cheese from raw goat's milk. *Food Microbiology*. 20: 575-582. [DOI: 10.1016/S0740-0020(02)00153-3]
- Rezac S., Kok C.R., Heermann M., Hutkins R. (2018). Fermented foods as a dietary source of live organisms. *Frontiers in Microbiology*. 9: 1785. [DOI: 10.3389/fmicb.2018.01785]
- Ribeiro A.C., Ribeiro S.D.A. (2010). Specialty products made from goat milk. *Small Ruminant Research*. 89: 225-233. [DOI: 10.1016/j.smallrumres.2009.12.048]
- Russo N., Caggia C., Pino A., Coque T.M., Arioli S., Randazzo C.L. (2018). *Enterococcus* spp. in Ragusano PDO and Pecorino Siciliano cheese types: a snapshot of their antibiotic resistance distribution. *Food and Chemical Toxicology*. 120: 277-286. [DOI: 10.1016/j.fct.2018.07.023]
- Scatassa M.L., Gaglio R., Cardamone C., Macaluso G., Arcuri L., Todaro M., Mancuso I. (2017). Anti-*Listeria* activity of lactic acid bacteria in two traditional Sicilian cheeses. *Italian Journal of Food Safety*. 6: 6191. [DOI: 10.4081/ijfs.2017.6191]
- Scatassa M.L., Gaglio R., Macaluso G., Francesca N., Randazzo W., Cardamone C., Di Grigoli A., Moschetti G., Settanni L. (2015). Transfer, composition and technological characterization of the lactic acid bacterial populations of the wooden vats used to produce traditional stretched cheeses. *Food Microbiology*. 52: 31-41. [DOI: 10.1016/j.fm.2015.06.008]
- Settanni L., Di Grigoli A., Tornambé G., Bellina V., Francesca N., Moschetti G., Bonanno A. (2012). Persistence of wild *Streptococcus thermophilus* strains on wooden vat and during the manufacture of a traditional Caciocavallo type cheese. *International Journal of Food Microbiology*. 155: 73-81. [DOI: 10.1016/j.ijfoodmicro.2012.01.022]
- Settanni L., Gaglio R., Guarcello R., Francesca N., Carpino S., Sannino C., Todaro M. (2013). Selected lactic acid bacteria as a hurdle to the microbial spoilage of cheese: application on a traditional raw ewes' milk cheese. *International Dairy Journal*. 32: 126-132. [DOI: 10.1016/j.idairyj.2013.04.010]
- Settanni L., Guarcello R., Gaglio R., Francesca N., Aleo A., Felis G.E., Moschetti G. (2014). Production, stability, gene sequencing and in situ anti-*Listeria* activity of mundticin KS expressed by three *Enterococcus mundtii* strains. *Food Control*. 35: 311-322. [DOI: 10.1016/j.foodcont.2013.07.022]
- Settanni L., Moschetti G. (2010). Non-starter lactic acid bacteria used to improve cheese quality and provide health benefits. *Food Microbiology*. 27: 691-697. [DOI: 10.1016/j.fm.2010.05.023]
- Settanni L., Moschetti G. (2014). New trends in technology and identity of traditional dairy and fermented meat production processes: preservation of typicality and hygiene. *Trends in Food Science and Technology*. 37: 51-58. [DOI: 10.1016/j.tifs.2014.02.006]
- Suzzi G., Caruso M., Gardini F., Lombardi A., Vannini L., Guertzoni M.E., Andrighetto C., Lanorte M.T. (2000). A survey of the enterococci isolated from an artisanal Italian goat's cheese (semicotto caprino). *Journal of Applied Microbiology*. 89: 267-274. [DOI: 10.1046/j.1365-2672.2000.01120.x]
- Turchi B., Nuvoloni R., Fratini F., Pedonese F., Ebani V.V., Cerri D. (2011). Caciotta Della Garfagnana cheese: selection and evaluation of autochthonous mesophilic lactic acid bacteria as starter cultures. *Italian Journal of Animal Science*. 10: e22. [DOI: 10.4081/ijas.2011.e22]
- Weisburg W.G., Barns S.M., Pelletier D.A., Lane D.J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology*. 173: 697-703. [DOI:10.1128/jb.173.2.697-703.1991]