



## Turkey (*Meleagris gallopavo f. domestica*) Meat as a Potential Reservoir for Dissemination of Antimicrobial Resistant and Shigatoxigenic *Escherichia coli* Strains

M. Bagheri <sup>1\*</sup>, R. Ghanbarpour <sup>2</sup>, Z. Amirpoorsaid <sup>1</sup>, P. Mohseni <sup>3</sup>, M. Hesam-Arefi <sup>3</sup>, M. Jajarmi <sup>3</sup>

1. Department of Food Science and Technology, Bardsir Faculty of Agriculture, Shahid Bahonar University of Kerman, Kerman, Iran

2. Molecular Microbiology Research Group, Shahid Bahonar University of Kerman, Kerman, Iran

3. Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran

### HIGHLIGHTS

- *Escherichia coli* was detected in 75.78% of turkey carcasses, with 52.7% of isolates being multidrug-resistant.
- A significant resistance to chloramphenicol (63.89%), nalidixic acid (59.72%), and florfenicol (56.94%) was detected.
- The virulence genes *stx1* (4.17%), *stx2* (1.39%), and *eae* (2.77%) were identified in the isolates.
- The *E. coli* isolates were assigned to phylotypes A, B1, C, D, and E, while 9.72% remained unclassified.
- The findings highlighted the public health risks associated with antibiotic-resistant STEC and EPEC strains in turkey meat.

### Article type

Original article

### Keywords

Shiga-Toxigenic *Escherichia coli*

Beta-Lactamase

Antibiotic Resistance

Phylogeny

Turkeys

Abattoirs

### Article history

Received: 17 Jul 2024

Revised: 30 Aug 2025

Accepted: 15 Sep 2025

### Abbreviations

AMR=Antimicrobial-Resistant

CLSI=Clinical and Laboratory

Standards Institute

EHEC=Enterohemorrhagic

*Escherichia coli*

EPEC=Enteropathogenic

*Escherichia coli*

ESBL=Extended-Spectrum Beta-

Lactamase

MDR=Multi-Drug Resistant

PCR=Polymerase Chain

Reaction

STEC=Shiga Toxin-Producing

*Escherichia Coli*

### ABSTRACT

**Background:** Enteropathogenic and Shiga toxin-producing *Escherichia coli* (*E. coli*) are the two main pathotypes capable of causing serious human infections, especially when resistant to antibiotics.

**Methods:** In this work, 95 turkey carcasses were swabbed in slaughterhouse over four months (June–September 2023). Antibiotic resistance was evaluated by a disk diffusion method named Kirby–Bauer against nine antimicrobial agents. Three virulence genes including *stx1*, *stx2*, *eae*, three resistance genes including *bla<sub>TEM</sub>*, *bla<sub>SHV</sub>*, *bla<sub>CTX-M</sub>*, and four phylogenetic markers (*arpA*, *chuA*, *yjaA*, *TspE4.C2*) were screened by Polymerase Chain Reaction (PCR) method. Data were analyzed using Microsoft Excel and SPSS (version 24), with the chi-square test at a significance level of  $p \leq 0.05$ .

**Results:** Out of 95 carcasses, 72 (75.78%) were *E. coli*-positive. Among the *E. coli* isolates, 63.89% were resistant to chloramphenicol, 59.72% to nalidixic acid, and 56.94% to florfenicol. One isolate, classified as extended-spectrum beta-lactamase positive, belonged to phylogroup D and showed simultaneous resistance to four antibiotics without harboring the resistance genes studied. Overall, 52.7% of the *E. coli* isolates were recognized as Multi-Drug Resistant (MDR). Profiles of resistance genes included *bla<sub>TEM</sub>* (23.61%), *bla<sub>TEM</sub>/bla<sub>SHV</sub>* (2.77%), and *bla<sub>CTX-M</sub>* (1.38%). Virulence genes were detected in six isolates: *stx1* (4.17%), *stx2* (1.39%), and *eae* (2.77%). Phylogenetic analysis revealed five groups: A (19.44%), B1 (36.11%), C (5.55%), D (13.8%), and E (15.27%), while 9.72% remained unclassified.

**Conclusion:** The occurrence of *E. coli* isolates harboring virulence and antibiotic-resistance genes in turkey carcasses underscores serious public health risks.

The significant frequency of Multi-Drug Resistant (MDR) *E. coli* isolates highlights the need for improved monitoring and control measures throughout the food chain.

© 2026, Shahid Sadoughi University of Medical Sciences. This is an open access article under the Creative Commons Attribution 4.0 International License.

\* Corresponding author (M. Bagheri)

E-mail: ma.bagheri@uk.ac.ir

ORCID ID: <https://orcid.org/0000-0003-0522-9747>

**To cite:** Bagheri, M., Ghanbarpour, R., Amirpoorsaid, Z., Mohseni, P., Hesam-Arefi, M. and Jajarmi, M. (2026) 'Turkey (*Meleagris gallopavo f. domestica*) Meat as a Potential Reservoir for Dissemination of Antimicrobial Resistant and Shigatoxigenic *Escherichia coli* Strains', *Journal of Food Quality and Hazards Control*, 13(1), pp. 40-49.

## Introduction

Turkey meat (*Meleagris gallopavo* f. *domestica*) is one of the most consumed meats worldwide, following chicken meat (Connolly and Campbell, 2023). Meat products of turkey may be one of the putative reservoirs for several key foodborne pathogens, such as *Escherichia coli*, *Staphylococcus* spp., *Salmonella* spp., *Clostridium* spp., *Campylobacter* spp., *Listeria* spp., and *Arcobacter* spp. (Silva, Vidal and Junior, 2017). *E. coli* is of particular significance due to its predominant presence within the gastrointestinal microbiota in warm-blooded animals (Foster-Nyarko and Pallen, 2022).

*E. coli* is mainly commensal, but some strains exhibit pathogenicity due to distinct virulence factors (Pakbin, Brück and Rossen, 2021). Among the different pathotypes used to classify pathogenic *E. coli* strains, Shiga toxin-producing (STEC) and enteropathogenic *E. coli* (EPEC) are regarded as the most relevant to public health (Pakbin, Brück and Rossen, 2021). STECs produce two types of Shiga toxins named Stx1 and Stx2, which are encoded by *stx1* and *stx2* genes, whereas a distinctive feature of EPEC is its ability to cause attaching and effacing lesions, mediated by the outer membrane protein intimin, encoded by the *eae* gene (Pakbin, Brück and Rossen, 2021). The co-occurrence of intimin and *stx1* and/or *stx2* in *E. coli* strains defines the clinically significant pathotype called Enterohemorrhagic *E. coli* (EHEC; Zarei et al., 2021). *Stx* is a major virulence factor produced by STEC and EHEC, contributing to HC (hemorrhagic colitis) and HUS (Hemolytic Uremic Syndrome) (Freedman, Van De Kar and Tarr, 2023). Intimate adherence of bacteria to gut epithelial cells, causes disruption of the microvilli and formation of pedestal-like structures (Mare et al., 2021; Sokolovic et al., 2022).

One of the major challenges in infections caused by EPEC, STEC, and EHEC is the presence of antibiotic resistance in *E. coli* strains (Bouzari et al., 2018). The emergence of Antimicrobial-Resistant (AMR) *E. coli* in some food-producing animals, including poultry, has been increasingly reported worldwide (Van Boeckel et al., 2019). Studies have identified AMR *E. coli* isolates in meat products of turkey and other poultry, raising concerns about their potential foodborne transmission to humans (Khaita et al., 2008). These AMR strains may be resistant to fluoroquinolones and cephalosporins, which are essential for the treatment of severe human infections (Shrestha et al., 2022). The global health implications of AMR *E. coli* are significant, as these strains not only compromise treatment efficacy but also serve as reservoirs of resistance genes that may disseminate to other pathogenic/non-pathogenic bacteria (Martinez, 2009). Factors contributing to the emergence of AMR *E. coli* include improper antibiotic use, administration without

prior antimicrobial susceptibility testing, and the utilization of counterfeit drugs; such practices exert selective pressure, favoring resistant bacteria while disadvantaging antibiotic-susceptible strains. Addressing this issue is crucial to public health, as it underscores the need for monitoring AMR *E. coli* prevalence in poultry and implementing more stringent antimicrobial stewardship practices (Endale, Mathewos and Abdeta, 2023).

*E. coli* populations are clonal and can be assigned to distinct phylogroups including A, B1, B2, D, C, E, F, G, and *Escherichia* cryptic clade I based on screening of four genetic markers: *arpA*, *TspE4.C2*, *yjaA* and *chuA* (Beghain et al., 2018). These phylogenetic groups reflect the virulence, evolutionary history, ecological niche and both genotypic/phenotypic characteristics of *E. coli* strains (Stoppe et al., 2017). Understanding the phylogroup is crucial, as certain groups are more commonly associated with virulence and antibiotic resistance features, thereby influencing the pathogenic potential and clinical outcomes of *E. coli* infections characteristics (Tenailon et al., 2010).

Therefore, this study aims to investigate the prevalence of virulence genes and antimicrobial resistance factors among various phylogenetic groups of *E. coli* isolates from turkey meat. The results of this study may shed light on the contribution of turkey meat to the epidemiology of virulent and AMR *E. coli* strains.

## Materials and methods

### Collection of samples and isolation of *E. coli*

In total, 95 swab samples were collected from the carcasses of 95 turkeys in the slaughterhouse after chilling during four months (June to September 2023), before any secondary processing; Samples were taken by swabbing the interior surface of the turkey carcasses at the end of the slaughter line, prior to carcass cutting and packaging. A sterile swab was used to thoroughly swab the internal cavity of each carcass, moving from one end to the other 5 to 10 times to ensure sufficient sample collection. The swab was then placed into a transport medium for subsequent microbiological analysis. This standardized approach was aimed to ensure consistency and reproducibility across all samples. The sample size was estimated based on the formula:

$$\text{Sample size} = \frac{Z_{1-\frac{\alpha}{2}}^2 \times p \times (1-p)}{d^2} = \frac{3.84 \times 0.4 \times 0.6}{0.01} = 92.16$$

Z value for the 95% confidence interval [error level ( $\alpha$ ): 0.05] = 1.96;  $Z^2 = 3.84$

$p$  [the projected proportion based on earlier research (Nwankwo et al., 2021)] = 0.4 (or 40%)

$1 - p = 0.6$  (or 60%)

$d$  (the error rate chosen by the investigators in this work) = 0.1 (or 10%);  $d^2 = 0.01$

The swab samples were collected using aseptic techniques, stored in sterile tubes containing transport medium (Cary Blair; Himedia, India) under controlled temperature conditions (4°C), and immediately handled to laboratory mostly within 2 h and sometimes up to the 12 h. The swabs were transferred to the laboratory within 12 h. The samples were streaked on MacConkey agar (Merck, Germany) and incubated at 37 °C (Memmert, Germany) for 24 h. The lactose-positive colonies were biochemically confirmed by IMViC biochemical tests, including indole, methyl red, Voges-Proskauer, and citrate as *E. coli* strains (Markey *et al.*, 2013). Then, one strain from each bird was considered for virulence, antimicrobial resistance, and phylogenetic assessments.

#### Phenotypic evaluation of the *E. coli* isolates

Antimicrobial resistance was determined in this work via the Kirby-Bauer disc diffusion method, performed in accordance with the Clinical and Laboratory Standards Institute (CLSI) guidelines (CLSI, 2021); the antibiotics used (Padtan Teb co., Iran) included ceftazidime, cefotaxime, cefotaxime clavulanate, ceftazidime clavulanate, florfenicol, chloramphenicol, enrofloxacin, nalidixic acid, and ciprofloxacin. The reference strain *E. coli* ATCC 25922 was used as the quality control strain, and the inhibition zone diameters were interpreted according to CLSI guidelines (CLSI, 2021). Isolates showing an increase of  $\geq 5$  mm in the inhibition zone for cefotaxime-clavulanate relative to cefotaxime alone, or for ceftazidime-clavulanate relative to ceftazidime alone, were considered Extended-Spectrum  $\beta$ -Lactamase-producing *E. coli* (ESBL; CLSI, 2021).

#### Polymerase Chain Reaction (PCR) for virulence, antimicrobial resistance, and phylogenetic genes

DNA extraction was carried out using boiling lysis method. In brief, *E. coli* colonies were suspended in 400  $\mu$ l water (distilled and sterile) and heated at 98-100 °C for 10 min in boiling water. The lysates were then placed on (10 min) and centrifuged (13,000  $\times$  g; 1 min). The supernatant, containing the released DNA, was collected and used as the template for subsequent reactions. DNA yield and purity were evaluated with NanoDrop (BioTek Epoch, USA) by measuring absorbance at 260/280 nm.

The occurrence of the genes including *stx*<sub>1</sub>, *stx*<sub>2</sub>, and *eae*

were screened (Paton and Paton, 2002) by PCR (Bio-Rad, USA). Sakai strain of *E. coli* (*eae*<sup>+</sup>, *stx*<sub>1</sub><sup>+</sup>, and *stx*<sub>2</sub><sup>+</sup>) was used as a control<sup>+</sup> while *E. coli* strain MG1655 (*eae*<sup>-</sup>, *stx*<sub>1</sub><sup>-</sup>, and *stx*<sub>2</sub><sup>-</sup>) was employed as a control<sup>-</sup>, sourced from Shahid Bahonar University of Kerman. Additionally, the *E. coli* strains were screened for *bla*<sub>CTX-M</sub>, *bla*<sub>TEM</sub>, and *bla*<sub>SHV</sub> as antibiotic resistance genes (Roschanski *et al.*, 2014) (Table 1).

Detection of four sequences (*arpA*, *chuA*, *yjaA*, and *TspE4.C2*) using PCR (multiplex) (Clermont *et al.*, 2013) led to phylogenetic grouping of *E. coli* into 7 groups and some clades. Isolates that were *arpA* positive but *chuA*, *yjaA*, and *TspE4.C2* negative were allocated to phylogroup A, whereas those that were *arpA* positive, *chuA* and *yjaA* negative, but *TspE4.C2* positive were classified as B1. Phylogroup B2 included isolates that were *arpA* negative, *chuA* positive, and positive for at least one of *yjaA* or *TspE4.C2*. Isolates with the profile *arpA* positive, *chuA* negative, *yjaA* positive, and *TspE4.C2* negative were subjected to C-specific PCR; if this reaction was positive, they were assigned to C, otherwise they were reclassified as A. Strains that were *arpA* positive and *chuA* positive but *yjaA* negative (with *TspE4.C2* either negative or positive) were tested using E-specific primers; isolates yielding a positive result were designated as E, and those testing negative were placed in D. Isolates showing *arpA* positive, *chuA* positive, *yjaA* positive, and *TspE4.C2* negative were similarly screened with E-specific primers: E-positive strains were classified as E, whereas E-negative ones were assigned to *Escherichia* cryptic clade I and confirmed using cryptic clade-specific primers. Strains that were *arpA* negative, *chuA* positive, *yjaA* negative, and *TspE4.C2* negative were grouped as F, and isolates displaying *arpA* negative, *chuA* negative, *yjaA* positive, and *TspE4.C2* negative were assigned to *Escherichia* cryptic clade I or II after confirmation with cryptic clade-specific primers (Table 1). Other genotypes were considered as unknown phylogroups.

A 1.5% agarose gel in a horizontal electrophoresis unit was used to resolve the PCR products at 100 V for 45 minutes (Cleaver Scientific, United Kingdom) and then visualized by fluorescence staining using Greenview dye (Parstous Biotechnology, Iran) and gel documentation system (Vilber Lourmat co., Quantum software version ST4, France). Amplicon sizes were determined using two types of DNA marker including 50 bp and 100 bp (Parstous Biotechnology, Iran) as a reference.

**Table 1:** Sequences of primers used in this study

Gene	Sequence (5'-3')	PCR condition	Product size(bp)	Reference
<i>stx1</i>	F-ATAAATCGCCATTTCGTTGACTAC R-AGAACGCCCACTGAGATCATC	35 cycles: 94°C (90 s), 60°C (90 s), 72°C (90 s)	348	(Paton and Paton, 2002)
<i>stx2</i>	F-GGCACTGTCTCTGAACTGCTCC R-TCGCCAGTTATCTGACATTCTG	35 cycles: 94°C (90 s), 60°C (90 s), 72°C (90 s)	584	(Paton and Paton, 2002)
<i>eae</i>	F-GACCCGGCACAAGCATAAGC R-CCACGTGCAGCAACAAGAGG	35 cycles: 94°C (90 s), 60°C (90 s), 72°C (90 s)	482	(Paton and Paton, 2002)
<i>arpA</i>	F-AACGCTATTTCGCCAGCTTGC R-TCTCCCCATACCGTACGCTA	30 cycles: 94°C (5 s), 59°C (20 s) and 1 cycle 72°C (5 min)	400	(Clermont et al., 2013)
<i>chuA</i>	F-ATGGTACCGGACGAACCAAC R-TGCCGCCAGTACCAAGACA	30 cycles: 94°C (5 s), 59°C (20 s) and 1 cycle 72°C (5 min)	288	(Clermont et al., 2013)
<i>yjaA</i>	F-CAAACGTGAAGTGTGAGGAG R-AATGCGTTCCCTCAACCTGTG	30 cycles: 94°C (5 s), 59°C (20 s) and 1 cycle 72°C (5 min)	211	(Clermont et al., 2013)
<i>TspE4.C2</i>	F-CACTATTCGTAAGGTCATCC R-AGTTTATCGCTGCGGGTCCG	30 cycles: 94°C (5 s), 59°C (20 s) and 1 cycle 72°C (5 min)	152	(Clermont et al., 2013)
<i>bla<sub>TEM</sub></i>	Fa -GCGGAACCCCTATTTG Rb -ACCAATGCTTAATCAGTGAG	35 cycles: 95°C (30 s), 52°C (30 s), 72°C (60 s)	963	(Roschanski et al., 2014)
<i>bla<sub>SHV</sub></i>	F-TTATCTCCCTGTTAGCCACC R-GATTTGCTGATTTCCGCTCGG	35 cycles: 95°C (30 s), 54°C (30 s), 72°C (30 s)	795	(Roschanski et al., 2014)
<i>bla<sub>CTX-M</sub></i>	F-CGATGTGCAGTACCAGTAA R-TTAGTGACCAGAATCAGCGG	35 cycles: 95°C (30 s), 60°C (30 s), 72°C (60 s)	585	(Roschanski et al., 2014)

### Statistical analysis

To minimize potential confounding factors and biases, sampling was carried out over a 16-week period (4 months). During each sampling session, 5 to 10 samples were collected, and carcasses were selected in an alternating pattern (every other carcass) to ensure randomness and reduce selection bias. For each isolate, information on phylogroup status and antimicrobial resistance was compiled in Microsoft Excel 2016 and subsequently analyzed in SPSS version 24 (IBM) to obtain prevalence percentages through descriptive statistics. The chi-square test was applied to assess differences in frequencies and associations between variables, and a *p*-value of  $\leq 0.05$  was taken as the threshold for statistical significance.

### Results and discussion

#### Prevalence of *E. coli* strains

*E. coli* was detected in 72 out of 95 turkey carcasses (75.78%). The presence of *E. coli* on turkey carcasses suggests shortcomings in slaughterhouse hygiene, including possible fecal cross-contamination or improper

carcass handling. Such contamination represents a potential public health risk. *E. coli* may contribute substantially to the spread of virulence factors and antimicrobial resistance determinants.

#### Prevalence of phenotypic antimicrobial resistance

Antimicrobial resistance to seven widely used antibiotics was investigated; these antibiotics belonged to three different classes and were tested using the disc diffusion method. For amphenicols, 46 (63.89%) isolates showed resistance against chloramphenicol, and 41 (56.94%) isolates were resistant to florfenicol. For quinolones, resistance against enrofloxacin, nalidixic acid, and ciprofloxacin was detected in 23 (31.94%), 43 (59.72%), and 17 (23.61%) isolates, respectively. The lowest level of resistance was identified to  $\beta$ -lactams ( $p \leq 0.05$ ); only four strains were resistant against cefotaxime (5.56%), one was ceftazidime-resistant (1.38%), and only one isolate was ESBL-positive (1.38%) (Table 2). No differences were significantly observed in the frequency of antibiotic resistance among the phylogenetic groups ( $p > 0.05$ ).

**Table 2:** Prevalence of antimicrobial resistance phenotype among *Escherichia coli* isolates

Antimicrobial family	Antimicrobial resistance phenotype	Number	Percentage	95% CI
$\beta$ -lactams	Cefotaxime	4	5.56	0.26-10.85%
	Ceftazidime	1	1.38	0-4.09%
	Extended-Spectrum Beta-Lactamase-positive	1	1.38	0-4.09%
Amphenicols	Florfenicol	41	56.94	45.51-68.38%
	Chloramphenicol	46	63.89	52.79-74.98%
Quinolones	Enrofloxacin	23	31.94	21.17-42.71%
	Nalidixic acid	43	59.72	48.39-71.05%
	Ciprofloxacin	17	23.61	13.80-33.42%

The elevated levels of resistance to chloramphenicol, nalidixic acid, and florfenicol found in this study are consistent with previous findings in *E. coli* strains recovered from turkey meat and intestinal contents in Iran (Gholami-Ahangaran *et al.*, 2021); furthermore, the low levels of resistance to cefotaxime and ceftazidime observed in our study agreed with those reported previously in Germany, (Kaesbohrer *et al.*, 2012), Poland (Wasyl *et al.*, 2013), and Iran (Mousavi, Rahimi and Shakerian, 2020).

Lowest resistance level was against the  $\beta$ -lactams. The rate of antibiotic resistance for cefotaxime in this study was lower than study in Egypt (AbdelRahman *et al.*, 2020; Moawad *et al.*, 2018; Younis, Awad, and Mohamed, 2017), China (Wang *et al.*, 2021), and India (Sebastian *et al.*, 2021); only one isolate of our study was ESBL-positive. The lower levels of resistance against  $\beta$ -lactams in turkey meat, compared to florfenicol and chloramphenicol, may reflect differences in antibiotic usage practices in turkey production.

Antibiotic usage in veterinary medicine is a widespread approach either as a preventive measure or as additives to promote growth in livestock (Vázquez-Villanueva *et al.*, 2023). Unfortunately, this has resulted in the appearance of bacterial strains that exhibit resistance to multiple antibiotics, including florfenicol, chloramphenicol, cefotaxime, and ceftazidime. The present study demonstrates a considerable presence of AMR strains, suggesting a lack of adherence to recommended guidelines

and an overuse of antibiotics.

Resistance patterns and their distribution among phylogroups are shown in Table 3 and Figure 1. Twenty antimicrobial resistance profiles were observed in which florfenicol-chloramphenicol was the most prevalent AMR profile (12 isolates; 16.67%). Thirty-eight (52.7%) isolates were recognized as Multi-Drug Resistant (MDR; a strain, resistant to at least one antibiotic from three different antimicrobial classes); this finding is comparable to the results reported from the USA (Davis *et al.*, 2018). Six strains were susceptible to all tested antimicrobials. The most common resistance pattern was related to FF, C profile (12 isolates, 16.69%). In Egypt, a work on *E. coli* strains isolated from turkey showed 100% of the isolates were MDR (Eid and Samir, 2019). The global dissemination of MDR strains as a consequence of excessive and uncontrolled use of antimicrobials has caused a serious problem, indicating the high prevalence of MDR isolates among poultry farms.

#### Detection of $\beta$ -lactamase genes


The most frequent antimicrobial resistance genotypic pattern was *bla*<sub>TEM</sub> (17 isolates; 23.61%), followed by *bla*<sub>TEM</sub>/*bla*<sub>SHV</sub> (2 isolates; 2.77%), and *bla*<sub>CTX-M</sub> (1 isolate; 1.38%). Fifty-two isolates (72.22%) possessed none of the screened antimicrobial resistance genes. The phylogenetic distribution pattern of the positive isolates is revealed in Table 4.

**Table 3:** Phenotypic antimicrobial resistance profiles and their distribution pattern among phylo-groups

Antimicrobial resistance profiles	Phylo-group (no.)						No. (%)
	A	B1	C	D	E	U	
C		1				2	3 (4.17)
FF	1	1		1	1		4 (5.57)
NA		1	2	3	1		7 (9.73)
CTX		1					1 (1.38)
FF, C	5	3			1	3	12 (16.69)
C, NA						1	1 (1.38)
FF, NA		1					1 (1.38)
C, CTX	1						1 (1.38)
NA, CIP			1				1 (1.38)
FF, CTX					1		1 (1.38)
NA, NFX		1			1		2 (2.77)
FF, C, NA		3		3	2		8 (11.12)
FF, C, CIP				1			1 (1.38)
C, NA, NFX		1					1 (1.38)
NA, NFX, CIP	1			1	1		3 (4.17)
FF, C, NA, CIP		1					1 (1.38)
FF, C, NA, NFX	2	3			1		6 (8.34)
C, NA, NFX, CIP	1	1		1*		1	4 (5.57)
C, NA, CTX, CAZ	1						1 (1.38)
FF, C, NA, NFX, CIP	2	4			1		7 (9.73)
Non-resistant isolates	1	4			1		6 (8.34)
Total (%)	15 (20.83)	26 (36.11)	3 (4.17)	10 (13.89)	11 (15.28)	7 (9.72)	72 (100)

\*This strain is the only Extended-Spectrum Beta-Lactamase-producing *Escherichia coli* isolate.

C=chloramphenicol; CAZ=ceftazidime; CIP=ciprofloxacin; CTX=Cefotaxime; FF=florfenicol; NA=nalidixic acid; NFX=enrofloxacin



**Figure 1:** Heat map of phenotypic antimicrobial resistance patterns, phylogenetic background, virulence genes, and antimicrobial resistance genes. C=chloramphenicol; CAZ=ceftazidime; CIP=ciprofloxacin; CTX=Cefotaxime; FF=florfenicol; NA=nalidixic acid; NFX=enrofloxacin

**Table 4:** Prevalence of antimicrobial resistance genotype and distribution pattern of phylogenetic groups among *Escherichia coli* isolates

Variables	Phylo-group						No. (%)	95% confidence interval
	A	B1	C	D	E	U		
<i>bla</i> <sub>TEM</sub>	7	3	1	1	4	1	17 (23.61)	13.8%-33.42%
<i>bla</i> <sub>TEM</sub> / <i>bla</i> <sub>SHV</sub>			1		1		2 (2.77)	0%-6.57%
<i>bla</i> <sub>CTX-M</sub>				1			1 (1.38)	0%-4.09%
No resistance gene	8	23	1	8	6	6	52 (72.22)	61.88%-82.57%
Total	15	26	3	10	11	7	72 (100)	-

Higher frequency of *bla* gene-positive strains in poultry and turkey has been reported in Egypt (AbdelRahman *et al.*, 2020), Ghana (Eibach *et al.*, 2018), Türkiye (Baran, Adıgüzel and Yüksel, 2020), and Brazil (Hoepers *et al.*, 2018). These studies are relevant as they demonstrate the widespread occurrence of *bla* genes in poultry across diverse geographic regions. The present study’s results on the *bla*<sub>CTX-M</sub> closely align with finding from Canada (Sheikh *et al.*, 2012) and Egypt (Moawad *et al.*, 2018).

It is also important to recognize the potential for horizontal transfer of ESBL genes to other bacteria. Such transfer events may promote the dissemination of resistance features to otherwise susceptible bacterial populations, thereby increasing the likelihood of therapeutic failure and making infection control more difficult.

*Prevalence of virulence genes*

In the current study, six strains carried at least one of the

three virulence genes under investigation; *stx*<sub>1</sub> (3 isolates; 4.17%), *stx*<sub>2</sub> (1 isolate; 1.39%) and *eae* (2 isolates; 2.77%). No virulence genes were detected in 66 isolates (91.67%). Details are presented in table 5.

Furthermore, six strains harbored at least one of the three virulence genes analyzed, which are known to be associated with STEC, EPEC, and EHEC pathotypes; the

prevalence of VG-positives was less than 5%. Comparable frequencies of *stx*<sub>1</sub>, *stx*<sub>2</sub>, and *eae* were found by Mousavi, Rahimi and Shakerian (2020) in Iran. However, some studies in Saudi Arabia (Hessain *et al.*, 2015), Austria (Mayrhofer *et al.*, 2004), and Canada (Bohaychuk *et al.*, 2006) did not detect virulence genes related to STEC, EPEC, and EHEC.

**Table 5:** Prevalence of virulence gene and distribution pattern of phylogenetic groups among *Escherichia coli* isolates

Variables	Phylo-group						No. (%)	95% confidence interval
	A	B1	C	D	E	U		
<i>eae</i>				1		1	2 (2.77)	0%-6.57%
<i>stx</i> <sub>1</sub>	1	1			1		3 (4.17)	0%-8.78%
<i>stx</i> <sub>2</sub>				1			1 (1.39)	0%-4.09%
No virulence gene	14	25	3	8	10	6	66 (91.67)	82.74%-96.88%
Total	15	26	3	10	11	7	72 (100)	-

The variation in sample size between studies may be one reason for the observed differences in findings. The lower prevalence of the virulence genes in turkey meat compared to other food sources such as beef can be attributed to biological and husbandry-related factors. It is well-established that ruminants, rather than birds, serve as the primary reservoirs of STEC. Therefore, the presence of these virulence genes is generally higher in ruminant-derived products. However, in traditional turkey farming systems, especially where turkeys are raised in close proximity to ruminants, the likelihood of STEC contamination may increase due to potential cross-contamination or environmental exposure.

Available evidence suggests that food animals might function as a reservoir for human infection. Also, retail meats could potentially serve as a vehicle for the transmission of these strains. Although the frequency of pathogenic *E. coli* isolates in turkey carcasses is low, their potential role in spreading virulent strains should not be overlooked, especially considering vulnerable populations such as immunocompromised individuals, for whom even minimal exposure may pose significant health risks.

#### Prevalence of phylogenetic groups

Totally, among the 72 strains, seven phylogroups including A, B1, B2, C, D, E, and F were identified, while some isolates remained unassigned and were categorized as unknown (U); from high to low prevalence, 26 isolates (36.11%) belonged to B1, 14 isolates (19.44%) to A, 11 (15.27%) isolates to E, 10 isolates (13.8%) to D, 4 isolates (5.55%) to C, and 7 isolates (9.72%) didn't belong to any of the phylogenetic groups (U) (Table 3-5). A heat map of phenotypic antimicrobial resistance patterns and associated phylogenetic background, virulence genes, and antimicrobial resistance genotypic profile has been provided in Figure 1.

The dominance of B1 in our study is consistent with another work in Türkiye (Baran, Adıgüzel and Yüksel,

2020). The most prevalent phylotypes in China (Wang *et al.*, 2021), Spain (Egea *et al.*, 2012), and Iran (Asadi *et al.*, 2018) have been related to phylotype A. Various variables, for example body mass, age, diet, species, sex, type of nutrition, types of pollution, environment (geographic location and climate), bacteria (pathogenicity and resistance), method, sample size, and study time, account for differences in phylotypes across studies.

Most positive strains for  $\beta$ -lactam were found in A and B1, the phylogroups known as commensal phylogroups of *E. coli*. Therefore, commensal strains may represent a potential threat to public health because of their role in the transmission of antibiotic resistance. among different hosts and pathogens. Strains classified within phylogroup D were all negative for virulence genes. This finding is interesting because, in many studies, D has been introduced as a phylogroup related to pathogenic strains of *E. coli*. Although B2 and D are traditionally considered more pathogenic in humans and other hosts, some studies have reported that strains belonging to these groups may exhibit lower virulence potential. This observation suggests that phylogenetic grouping alone may not fully predict the pathogenicity of *E. coli* bacterium.

Phylogroups of *E. coli* may be associated with distinct antibiotic resistance patterns; however, this relationship is complex and influenced by multiple factors. The transfer of genes encoding resistance factors through mobile genetic elements can modify resistance patterns among different phylogroups (Xie, Ogura and Suzuki, 2022).

#### Conclusion

The findings of this study indicate a high prevalence of *E. coli* in turkey carcasses. turkey carcasses, with a substantial proportion of isolates identified as multidrug-resistant and harboring the genetic determinants related to virulence and antibiotic resistance. Occurrence of virulent and AMR *E. coli* isolates from turkey meat highlights

significant public health concerns, particularly in relation to food safety and antibiotic stewardship.

The results highlight the attention to regular surveillance of antimicrobial resistance patterns in poultry meat, as well as the implementation of effective hygiene practices throughout the slaughtering process. Improved slaughterhouse hygiene, personnel training, and adherence to international standards, including Hazard Analysis and Critical Control Points (HACCP), are essential for minimizing microbial contamination.

Given the limitations of the present study, small sample size and the selection of only one isolate per bird, further research is recommended to better capture the heterogeneity and dynamics of *E. coli* bacterium in poultry.

Future studies should involve larger and geographically diverse sampling, comparison with other poultry species, and an expanded panel of virulence genes to gain a better understanding of the pathogenic ability and antimicrobial resistance profiles of *E. coli* in turkey meat.

#### Author contributions

M.B. conceptualized and administrated the project and wrote, reviewed and edited the manuscript; R.G. supervised the research and reviewed the manuscript; Z.A., P.M., M.H.A., and M.J. performed the sampling, laboratory procedures, and data analysis. All authors read and approved the final manuscript.

#### Acknowledgments

The authors gratefully acknowledge the support of the manager and staff of Bardsir Industrial Poultry Slaughterhouse for their assistance in sampling for the present projects.

#### Conflicts of interest

The authors declare that there is no conflict of interest.

#### Funding

This research received no specific grant from any funding agency in the public, commercial, or non-profit sectors.

#### Ethical consideration

Not applicable.

#### References

AbdelRahman, M.A.A., Roshdy, H., Samir, A.H. and Hamed, E.A. (2020) 'Antibiotic resistance and extended-spectrum  $\beta$ -lactamase in *Escherichia coli* isolates from imported 1-day-old chicks, ducklings, and turkey poults', *Veterinary World*, 13(6), pp. 1037-1044. Available at: <https://doi.org/10.14202/vetworld.2020.1037-1044>

- Asadi, A., Salehi, T.Z., Jamshidian, M. and Ghanbarpour, R. (2018) 'ECOR phylotyping and determination of virulence genes in *Escherichia coli* isolates from pathological conditions of broiler chickens in poultry slaughter-houses of southeast of Iran', *Veterinary Research Forum*, 9(3), pp. 211-216. Available at: <https://doi.org/10.30466/vrf.2018.30827>
- Baran, A., Adigüzel, M. and Yüksel, M. (2020) 'Prevalence of antibiotic-resistant and extended-spectrum beta-lactamase-producing *Escherichia coli* in chicken meat from eastern Turkey', *Pakistan Veterinary Journal*, 40(3), pp. 355-359. Available at: <https://doi.org/10.29261/pakvetj/2020.047>
- Beghain, J., Bridier-Nahmias, A., Le Nagard, H., Denamur, E. and Clermont, O. (2018) 'ClermontTyping: an easy-to-use and accurate in silico method for *Escherichia coli* strain phylotyping', *Microbial Genomics*, 4(7), p. e000192. Available at: <https://doi.org/10.1099/mgen.0.000192>
- Bohaychuk, V.M., Gensler, G.E., King, R.K., Manninen, K.I., Sorensen, O., Wu, J.T., Stiles, M.E. and McMullen, L.M. (2006) 'Occurrence of pathogens in raw and ready-to-eat meat and poultry products collected from the retail marketplace in Edmonton, Alberta, Canada', *Journal of Food Protection*, 69(9), pp. 2176-2182. Available at: <https://doi.org/10.4315/0362-028X-69.9.2176>
- Bouzari, S., Farhang, E., Hosseini, S.M. and Alikhani, M.Y. (2018) 'Prevalence and antimicrobial resistance of Shiga toxin-producing *Escherichia coli* and enteropathogenic *Escherichia coli* isolated from patients with acute diarrhea', *Iranian Journal of Microbiology*, 10(3), pp. 151-157.
- Clermont, O., Christenson, J.K., Denamur, E. and Gordon, D.M. (2013) 'The Clermont *Escherichia coli* phylo-typing method revisited: Improvement of specificity and detection of new phylo-groups', *Environmental Microbiology Reports*, 5(1), pp. 58-65. Available at: <https://doi.org/10.1111/1758-2229.12019>
- Clinical and Laboratory Standards Institute (CLSI) (2021) *Performance Standards for Antimicrobial Susceptibility Testing*. 31st edn. CLSI supplement M100. Wayne, Pennsylvania, USA: Clinical and Laboratory Standards Institute.
- Connolly, G. and Campbell, W.W. (2023) 'Poultry consumption and human cardiometabolic health-related outcomes: A narrative review', *Nutrients*, 15(16), p. 3550. Available at: <https://doi.org/10.3390/nu15163550>
- Davis, G.S., Waits, K., Nordstrom, L., Grande, H., Weaver, B., Papp, K., Horwinski, J., Koch, B., Hungate, B.A., Liu, C.M. and Price, L.B. (2018) 'Antibiotic-resistant *Escherichia coli* from retail poultry meat with different antibiotic use claims', *BMC Microbiology*, 18, p. 174. Available at: <https://doi.org/10.1186/s12866-018-1322-5>
- Egea, P., López-Cerero, L., Torres, E., Del Carmen Gómez-Sánchez, M., Serrano, L., Sánchez-Ortiz, M.D.N., Rodríguez-Baño, J. and Pascual, A. (2012) 'Increased raw poultry meat colonization by extended spectrum beta-lactamase-producing *Escherichia coli* in the south of Spain', *International Journal of Food Microbiology*, 159(2), pp. 69-73. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2012.08.002>
- Eibach, D., Dekker, D., Boahen, K.G., Akenten, C.W., Sarpong, N., Campos, C.B., Berneking, L., Aepfelbacher, M., Krumkamp, R., Owusu-Dabo, E. and May, J. (2018) 'Extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in local and imported poultry meat in Ghana', *Veterinary Microbiology*, 217, pp. 7-12. Available at: <https://doi.org/10.1016/j.vetmic.2018.02.023>

- Eid, S. and Samir, A.H. (2019) 'Extended-spectrum beta-lactamase and Class I integrons in multidrug-resistant *Escherichia coli* isolated from turkeys', *Veterinary World*, 12(7), pp. 1167–1174. Available at: <https://doi.org/10.14202/vetworld.2019.1167-1174>
- Endale, H., Mathewos, M. and Abdeta, D. (2023) 'Potential causes of spread of antimicrobial resistance and preventive measures in one health perspective-a review', *Infection and Drug Resistance*, 16, pp. 7515–7545. Available at: <https://doi.org/10.2147/IDR.S428837>
- Foster-Nyarko, E. and Pallen, M.J. (2022) 'The microbial ecology of *Escherichia coli* in the vertebrate gut', *FEMS Microbiology Reviews*, 46(3), p. fuac008. Available at: <https://doi.org/10.1093/femsre/fuac008>
- Freedman, S.B., Van De Kar, N.C.A.J. and Tarr, P.I. (2023) 'Shiga toxin-producing *Escherichia coli* and the hemolytic-uremic syndrome', *New England Journal of Medicine*, 389(15), pp. 1402–1414. Available at: <https://doi.org/10.1056/NEJMra2108739>
- Gholami-Ahangaran, M., Moravvej, A.H., Safizadeh, Z., Sadeghi Nagoorani, V., Zokaei, M. and Ghasemian, S.O. (2021) 'The evaluation of ESBL genes and antibiotic resistance rate in *Escherichia coli* strains isolated from meat and intestinal contents of turkey in Isfahan, Iran', *Iranian Journal of Veterinary Research*, 22(4), pp. 318–325. Available at: <https://doi.org/10.22099/ijvr.2021.39493.5737>
- Hessain, A.M., Al-Arfaj, A.A., Zakri, A.M., El-Jakee, J.K., Al-Zogibi, O.G., Hemeg, H.A. and Ibrahim, I.M. (2015) 'Molecular characterization of *Escherichia coli* O157: H7 recovered from meat and meat products relevant to human health in Riyadh, Saudi Arabia', *Saudi Journal of Biological Sciences*, 22(6), pp. 725–729. Available at: <https://doi.org/10.1016/j.sjbs.2015.06.009>
- Hoepers, P.G., Silva, P.L., Rossi, D.A., Valadares Júnior, E.C., Ferreira, B.C., Zuffo, J.P., Koerich, P.K. and Fonseca, B.B. (2018) 'The association between extended spectrum beta-lactamase (ESBL) and ampicillin C (AmpC) beta-lactamase genes with multidrug resistance in *Escherichia coli* isolates recovered from turkeys in Brazil', *British Poultry Science*, 59(4), pp. 396–401. Available at: <https://doi.org/10.1080/00071668.2018.1468070>
- Kaesbohrer, A., Schroeter, A., Tenhagen, B.-A., Alt, K., Guerra, B. and Appel, B. (2012) 'Emerging antimicrobial resistance in commensal *Escherichia coli* with public health relevance', *Zoonoses and Public Health*, 59, pp. 158–165. Available at: <https://doi.org/10.1111/j.1863-2378.2011.01451.x>
- Khaita, M.L., Oloya, J., Doetkott, D. and Kegode, R. (2008) 'Antimicrobial resistance and association with class I integrons in *Escherichia coli* isolated from turkey meat products', *Journal of Food Protection*, 71(8), pp. 1679–1684. Available at: <https://doi.org/10.4315/0362-028X-71.8.1679>
- Mare, A.D., Ciurea, C.N., Man, A., Tudor, B., Moldovan, V., Decean, L. and Toma, F. (2021) 'Enteropathogenic *Escherichia coli*—a Summary of the literature', *Gastroenterology Insights*, 12(1), pp. 28–40. Available at: <https://doi.org/10.3390/gastroent12010004>
- Markey, B., Leonard, F., Archambault, M., Cullinane, A. and Maguire, D. (2013) *Clinical Veterinary Microbiology*. 2<sup>nd</sup> edn. London, UK: Elsevier Health Sciences.
- Martinez, J.L. (2009) 'The role of natural environments in the evolution of resistance traits in pathogenic bacteria', *Proceedings of the Royal Society B: Biological Sciences*, 276(1667), pp. 2521–2530. Available at: <https://doi.org/10.1098/rspb.2009.0320>
- Mayrhofer, S., Paulsen, P., Smulders, F.J.M. and Hilbert, F. (2004) 'Antimicrobial resistance profile of five major food-borne pathogens isolated from beef, pork and poultry', *International Journal of Food Microbiology*, 97(1), pp. 23–29. Available at: <https://doi.org/10.1016/j.ijfoodmicro.2004.04.006>
- Moawad, A.A., Hotzel, H., Neubauer, H., Ehrlich, R., Monecke, S., Tomaso, H., Hafez, H.M., Roesler, U. and El-Adawy, H. (2018) 'Antimicrobial resistance in Enterobacteriaceae from healthy broilers in Egypt: emergence of colistin-resistant and extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli*', *Gut Pathogens*, 10, p. 39. Available at: <https://doi.org/10.1186/s13099-018-0266-5>
- Mousavi, R., Rahimi, E. and Shakerian, A. (2020) 'Incidence and profiles of antibiotic resistance and virulence markers of the *Escherichia coli* O157 bacteria recovered from poultry meat', *Egyptian Journal of Veterinary Sciences*, 51(2), pp. 215–223. Available at: <https://doi.org/10.21608/ejvs.2020.20516.1141>
- Nwankwo, I.O., Ezenduka, E.V., Nwanta, J.A., Ogugua, A.J. and Audu, B.J. (2021) 'Prevalence of *Campylobacter* spp. and antibiotics resistant *E. coli* on poultry carcasses and handlers' hands at Ikpa slaughter, Nsukka, Nigeria', *Notulae Scientia Biologicae*, 13(2), p. 10866. Available at: <https://doi.org/10.15835/nsb13210866>
- Pakbin, B., Brück, W.M. and Rossen, J.W.A. (2021) 'Virulence factors of enteric pathogenic *Escherichia coli*: a review', *International Journal of Molecular Sciences*, 22(18), p. 9922. Available at: <https://doi.org/10.3390/ijms22189922>
- Paton, A.W. and Paton, J.C. (2002) 'Direct detection and characterization of Shiga toxin-producing *Escherichia coli* by multiplex PCR for stx1, stx2, eae, ehxA, and saa', *Journal of Clinical Microbiology*, 40(1), pp. 271–274. Available at: <https://doi.org/10.1128/jcm.40.1.271-274.2002>
- Roschanski, N., Fischer, J., Guerra, B. and Roesler, U. (2014) 'Development of a multiplex real-time PCR for the rapid detection of the predominant beta-lactamase genes CTX-M, SHV, TEM and CIT-type AmpCs in Enterobacteriaceae', *PLoS One*, 9(7), p. e100956. Available at: <https://doi.org/10.1371/journal.pone.0100956>
- Sebastian, S., Tom, A.A., Babu, J.A. and Joshy, M. (2021) 'Antibiotic resistance in *Escherichia coli* isolates from poultry environment and UTI patients in Kerala, India: a comparison study', *Comparative Immunology, Microbiology and Infectious Diseases*, 75, p. 101614. Available at: <https://doi.org/10.1016/j.cimid.2021.101614>
- Sheikh, A.A., Checkley, S., Avery, B., Chalmers, G., Bohaychuk, V., Boerlin, P., Reid-Smith, R. and Aslam, M. (2012) 'Antimicrobial resistance and resistance genes in *Escherichia coli* isolated from retail meat purchased in Alberta, Canada', *Foodborne Pathogens and Disease*, 9(7), pp. 625–631. Available at: <https://doi.org/10.1089/fpd.2011.1078>
- Shrestha, R.D., Agunos, A., Gow, S.P., Deckert, A.E. and Varga, C. (2022) 'Associations between antimicrobial resistance in fecal *Escherichia coli* isolates and antimicrobial use in Canadian turkey flocks', *Frontiers in Microbiology*, 13, p. 954123. Available at: <https://doi.org/10.3389/fmicb.2022.954123>
- Silva, H.O., Vidal, A.M.C. and Junior, O.D.R. (2017) 'Pathogenic bacteria in turkey meat: A review', *Revista Brasileira de Higiene e Sanidade Animal*, 11(3), pp. 338–353.
- Sokolović, M., Šimpraga, B., Amšel-Zelenika, T., Berendika, M. and Krstulović, F. (2022) 'Prevalence and characterization of Shiga toxin producing *Escherichia coli* isolated from animal feed in Croatia', *Microorganisms*, 10(9), p. 1839. Available at: <https://doi.org/10.3390/microorganisms10091839>

- Stoppe, N.D.C., Silva, J.S., Carlos, C., Sato, M.I.Z., Saraiva, A.M., Ottoboni, L.M.M. and Torres, T.T. (2017) 'Worldwide phylogenetic group patterns of *Escherichia coli* from commensal human and wastewater treatment plant isolates', *Frontiers in Microbiology*, 8, p. 2512. Available at: <https://doi.org/10.3389/fmicb.2017.02512>
- Tenaillon, O., Skumik, D., Picard, B. and Denamur, E. (2010) 'The population genetics of commensal *Escherichia coli*', *Nature Reviews Microbiology*, 8, pp. 207–217. Available at: <https://doi.org/10.1038/nrmicro2298>
- Van Boeckel, T.P., Pires, J., Silvester, R., Zhao, C., Song, J., Criscuolo, N.G., Gilbert, M., Bonhoeffer, S. and Laxminarayan, R. (2019) 'Global trends in antimicrobial resistance in animals in low-and middle-income countries', *Science*, 365(6459), p. eaaw1944. Available at: <https://doi.org/10.1126/science.aaw1944>
- Vázquez-Villanueva, J., Vázquez, K., Martínez-Vázquez, A.V., Wong-González, A., Hernández-Escareño, J., Cabrero-Martínez, O., Cruz-Pulido, W.L., Guerrero, A., Rivera, G. and Bocanegra-García, V. (2023) 'Molecular and antimicrobial susceptibility characterization of *Escherichia coli* isolates from bovine slaughterhouse process', *Antibiotics*, 12(2), p. 291. Available at: <https://doi.org/10.3390/antibiotics12020291>
- Wang, M., Jiang, M., Wang, Z., Chen, R., Zhuge, X. and Dai, J. (2021) 'Characterization of antimicrobial resistance in chicken source phylogroup F *Escherichia coli*: similar populations and resistance spectrums between *E. coli* recovered from chicken colibacillosis tissues and retail raw meats in Eastern China', *Poultry Science*, 100(9), p. 101370. Available at: <https://doi.org/10.1016/j.psj.2021.101370>
- Wasył, D., Hoszowski, A., Zając, M. and Szulowski, K. (2013) 'Antimicrobial resistance in commensal *Escherichia coli* isolated from animals at slaughter', *Frontiers in Microbiology*, 4, p. 221. Available at: <https://doi.org/10.3389/fmicb.2013.00221>
- Xie, H., Ogura, Y. and Suzuki, Y. (2022) 'Persistence of antibiotic-resistant *Escherichia coli* strains belonging to the B2 phylogroup in municipal wastewater under aerobic conditions', *Antibiotics*, 11(2), p. 202. Available at: <https://doi.org/10.3390/antibiotics11020202>
- Younis, G., Awad, A. and Mohamed, N. (2017) 'Phenotypic and genotypic characterization of antimicrobial susceptibility of avian pathogenic *Escherichia coli* isolated from broiler chickens', *Veterinary World*, 10(10), pp. 1167–1172. Available at: <https://doi.org/10.14202/vetworld.2017.1167-1172>
- Zarei, O., Shokoohzadeh, L., Hossainpour, H. and Alikhani, M.Y. (2021) 'The prevalence of Shiga toxin-producing *Escherichia coli* and enteropathogenic *Escherichia coli* isolated from raw chicken meat samples', *International Journal of Microbiology*, 2021, p. 3333240. Available at: <https://doi.org/10.1155/2021/3333240>