



Contamination of *Cryptosporidium* spp. Oocysts in Raw Vegetables Produced in Koya City, Iraq

H. Sleman Ali^{1,2}, S.N. Mageed³, Gh.R. Jahed Khaniki^{1*}✉, N. Shariatifar¹, M. Yunesian¹, M. Rezaeian⁴, K.Kh. Saleh²

1. Division of Food Safety and Hygiene, Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Health Prevention, Koya Technical Institute, Koya City-Erbil, Iraq

3. Department of Medical Microbiology, Koya University, University Park, Koya City-Erbil, Iraq

4. Department of Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

HIGHLIGHTS

- The average prevalence rate of *Cryptosporidium* was 6.5% in vegetables of Koya, Iraq.
- Contamination rates were not related to vegetable types and also sampling seasons.
- Serious control measures and training must be carried out in the studied area.

Article type

Original article

Keywords

Cryptosporidium
Prevalence
Vegetables
Iraq

Article history

Received: 1 Oct 2017

Revised: 19 Jan 2018

Accepted: 19 Feb 2018

ABSTRACT

Background: *Cryptosporidium* spp. is one of the most important parasitic pathogen which causes acute gastroenteritis in human. This study aimed to determine and compare the contamination rates of *Cryptosporidium* oocysts in some common vegetables that are cultured and consumed in Koya City, Iraq.

Methods: Totally, 400 vegetable samples were collected randomly from vegetable farms in Koya city during spring and summer, 2016. The vegetables, including celery, parsley, leek, radish, and green onion were examined for the presence of *Cryptosporidium* spp. by Ziehl-Neelsen acid-fast staining method. Data were analyzed by SPSS statistical software (version 21).

Results: The prevalence rate of *Cryptosporidium* spp. oocysts in parsley, celery, leek, radish, and green onion were 6.25, 7.50, 6.25, 6.25, and 6.25%, respectively, showing no significant differences ($p>0.05$). Also, *Cryptosporidium* spp. oocysts were detected in 12 out of 200 (6%) samples from spring and 14 out of 200 (7%) samples from summer, without any significant difference ($p>0.05$).

Conclusion: Our findings highlighted that if the proper washing as well as disinfecting actions are neglected before consumption of the vegetables, consumption of raw vegetables could be potentially hazardous, inducing *Cryptosporidium* infection in Iraqi people.

© 2018, Shahid Sadoughi University of Medical Sciences. This is an open access article under the Creative Commons Attribution 4.0 International License.

Introduction

Consumption of raw vegetables without proper washing is one of the most significant causes of human parasitic diseases (Said, 2012). *Cryptosporidium* spp. is an important zoonotic parasitic agent which its excreted

oocyst in the feces can infect the hosts, especially mammals. Although the oocysts cannot multiply out of the host but they have the ability to tolerate environmental adverse conditions for a several months until their inges-

* Corresponding author. ✉ ghjahed@sina.tums.ac.ir

To cite: Sleman Ali H., Mageed S.N., Jahed Khaniki Gh.R., Shariatifar N., Yunesian M., Rezaeian M., Saleh K.Kh. (2018). Contamination of *Cryptosporidium* spp. oocysts in raw vegetables produced in Koya city, Iraq. *Journal of Food Quality and Hazards Control*. 5: 89-93.

tion by a new suitable host. Within the new host, the parasite's life cycle begins again, involving multiplication through utilizing resources of infected host (Chalmers and Katzer, 2013; Checkley et al., 2015; Rossle and Latif, 2013).

The main clinical feature of cryptosporidiosis is watery diarrhea that can differ from relatively mild to quite severe. There is also complained of abdominal pain and mild fever by infected persons. The immunodeficiency persons are more sensitive to infection (Petri et al., 2000). This protozoan parasite can also spread by water and soil. Human get infection by fecal-oral transmission and consumption of raw vegetables contaminated by oocysts. The food handlers in catering or restaurants with poor personal hygiene and unsanitary practice are the major sources of the infection (Chalmers and Katzer, 2013). Irrigation of vegetables using untreated or fecal-contaminated water is the main contamination source of these products prior to harvesting (Al-Binali et al., 2006; Amoah et al., 2007; Pires et al., 2012); however, contamination could be occurred at the time of harvesting, transportation, handling, distribution, or even at home settings (Said, 2012; Tefera et al., 2014).

There are some studies performed previously to assess the fact of intestinal parasite transmission through raw vegetables in several countries such as Egypt (Said, 2012), Libya (Abougrain et al., 2010), Canada (Dixon et al., 2013), Ghana (Duedu et al., 2014), Saudi Arabia (Al-Megrm, 2010), Peru (Ortega et al., 1997), Philippines (Su et al., 2012), Nigeria (Damen et al., 2007), Iran (Rostami et al., 2016), and also Iraq (Hadi, 2011). In our literature study, we did not find any published data about the contamination of *Cryptosporidium* oocyst in produced edible raw vegetables in Koya city, Kurdistan Region-Iraq. Therefore, this study aimed to determine and compare the contamination rates of *Cryptosporidium* spp. oocysts in some common vegetables that are cultured and consumed in this area from Iraq.

Materials and methods

The samples

The Koy Sanjaq or Koya city is located in the Erbil Governorate of Iraqi-Kurdistan, close to the Iranian border. There are 27 agriculture farms which they cultivate and supply the raw vegetables. Totally, 400 vegetable samples were collected randomly from farms in Koya city during spring and summer, 2016. Eighty samples were collected from each of the most common raw consumed types of fresh vegetables, including celery, parsley, leek, radish, and green onion (Figure 1).

Parasitological examination

The vegetable samples were about 400-500 g from each vegetable type. Each sample was soaked in one L of physiological normal saline (0.95% NaCl) and followed by mechanical shaking for 15 min. The top layer was discarded and the remaining wash solution was filtered through a sieve (micron pore size) to remove large debris, and then centrifuged at 2000 rpm for 15 min. The supernatant was decanted, and after that a few drops of the sediment were put on the glass slides and analyzed for parasites using Ziehl-Neelsen acid-fast staining method. For this work, the smear on a slide prepared from the sediment; then, the smears were air dried, and then fixed by ethanol. In the next step, alkaline fuchsin was poured on each slide and heated, but not boiled, for 5 min. Each slide was then washed carefully by water and decolorized using sulfuric acid (2.5%) for about 1 min. Next, each slide was stained with 1% methylene blue for 1 min, rinsed with tap water, and air dried. The detection of *Cryptosporidium* spp. oocysts was carried out by light microscope observation with $\times 100$ oil-immersion objective lenses.

Data analysis

The collected data were analyzed by SPSS statistical software (version 21). The p value less than 0.05 were used to declare association.

Results

The average contamination rate of *Cryptosporidium* spp. oocysts in produced edible green vegetables in Koya city, Iraq was 6.5%. The results revealed that the prevalence rate of *Cryptosporidium* oocysts in parsley, celery, leek, radish, and green onion were 5 of 80 (6.25%), 6 of 80 (7.50%), 5 of 80 (6.25%), 5 of 80 (6.25%), and 5 of 80 (6.25%), respectively. There was no statistically significance difference between the vegetable types and the contamination rate of *Cryptosporidium* spp. oocysts ($p > 0.05$).

Cryptosporidium spp. oocysts were detected in 12 out of 200 (6%) samples from spring and 14 out of 200 (7%) samples from summer. However, no significant ($p > 0.05$) seasonal variation was found regarding contamination rates of *Cryptosporidium* spp. oocysts in the vegetables samples during spring and summer.

Discussion

In different areas of the world, the vegetables may be contaminated during pre-harvesting, harvesting, and post



Figure 1: The most common local vegetables in Koya City, Iraq sampled in this research

harvesting. Sometimes, untreated wastewater is used to irrigate the vegetables, therefore consumption of raw vegetables induce parasitic infection if no appropriate washing is done (Damen et al., 2007; Gupta et al., 2009; Insulander et al., 2008; Ma et al., 2014; Pires et al., 2012; Said, 2012). In our studied area in Iraq, most vegetable

farms are irrigated by wastewater and fertilized by natural fecal sources; these are likely the major causes of *Cryptosporidium* contamination rate of 6.5% found in the samples. Similar findings have been previously indicated by Al-Binali et al. (2006) from Saudi Arabia that reported the prevalence of 8.33 and 2.78% for *Cryptosporidi-*

um oocysts in watercress and leek samples, respectively. In another research, Dixon et al. (2013) found *Cryptosporidium* spp. in 5.9% of pre-cut salads and leafy greens sold in Ontario, Canada. In agreement with the results of the current survey, Ranjbar-Bahadori et al. (2013) showed that 6.6% of examined vegetable samples of Tehran, Iran were contaminated with *Cryptosporidium* spp. Also, they found green onion as the most contaminated vegetable among all vegetables samples. In addition, Ahmad et al. (2016) determined the occurrence of *Cryptosporidium* spp. oocyst in Egyptian raw vegetable salads, including onion (1.5%), dill and parsley (3.4%), tomato (1.1%), and carrot (2.4%) which in accordance with the present survey. However, much higher incidence rate of this parasite reported by some Egyptian researchers who found *Cryptosporidium* oocyst in 33.3% of parsley (Sherbini et al., 2016) and 19% of green onion samples (Said, 2012). Such high rate of *Cryptosporidium* was also seen in vegetables samples of Ilam, West of Iran reported as 30.7% (Avazpoor et al., 2015). The observed differences might be due to differences in environmental conditions, sanitary practice, and the sample size difference between the studies.

Based on the present research, contamination rates were not significantly related to vegetable type which is similar to the results of Maikai et al. (2013) who studied the prevalence of *Cryptosporidium* spp. in 200 vegetable samples marketed in Zaria metropolis, Nigeria with no difference between contamination rates in lettuce (48%), spinach (40%), and waterleaf (36%). Accordingly, Rahman et al. (2014) have reported that the rates of *Cryptosporidium* oocysts did not differ among the total examined vegetables in Bangladesh.

The contamination rates of *Cryptosporidium* spp. in our vegetable samples had no difference between spring and summer. Rostami et al. (2016) announced that parasitic contamination of Iranian vegetable samples was higher in warm seasons than in cold seasons. Similarly, Fallah et al. (2012) stated that although the rate of parasitic contamination rates in vegetable samples were not significantly different between spring and summer (as warm seasons), but this rates were reduced considerably in autumn and in winter (as cold seasons). Since our vegetable samples just were obtained in warm seasons, we could not achieve comprehensive comparison in this regards, which is limitation of this study.

Conclusion

Our findings highlighted that if the proper washing as well as disinfecting actions are neglected before consumption of the vegetables, the consumption of raw vegetables could be potentially hazardous for inducing

Cryptosporidium spp. infection in Iraqi people. Since inappropriate hygienic conditions and poor food handling practices are common in Iraq, serious control measures must be carried out by health authorities. Also, training of the farmers, food handlers and consumers is too important to improve their knowledge about risk of this parasitic infection and its preventive actions.

Author contributions

H.S.A. designed the study and wrote the manuscript. G.R.J.K. and S.N.M. conducted the study and supervised the analysis and the manuscript. N.S., M.Y., and M.R. drafted and consulted the analysis. H.S.A. and K.K.S. performed the experiments. All authors revised and approved the final manuscript.

Conflicts of interest

There is no conflict of interest.

Acknowledgements

This work was a part of funded MSc thesis of Hemn Sleman Ali, a student at Tehran University of Medical Sciences, International Campus. The authors would like to thank International Campus, Tehran University of Medical Sciences for providing financial support for this research (Grant No: 103-32910). This study was ethically approved by the local institutional review board.

References

- Abougrain A.K., Nahaisi M.H., Madi N.S., Saied M.M., Ghenghesh K.S. (2010). Parasitological contamination in salad vegetables in Tripoli-Libya. *Food Control*. 21: 760-762.
- Ahmad S.O., El Fadaly H.A., Zaki M.S., Barakat A.M.A. (2016). Incidence of zoonotic parasites in Egyptian raw vegetable salads. *Life Science Journal*. 13: 27-31.
- Al-Binali A.M., Bello C.S., El-Shewy K., Abdulla S.E. (2006). The prevalence of parasites in commonly used leafy vegetables in South Western Saudi Arabia. *Saudi Medical Journal*. 27: 613-616.
- Al-Megrm W.I. (2010). Prevalence of intestinal parasites in leafy vegetables in Riyadh, Saudi Arabia. *International Journal of Tropical Medicine*. 5: 20-23.
- Amoah P., Drechsel P., Abaidoo R.C., Klutse A. (2007). Effectiveness of common and improved sanitary washing methods in selected cities of West Africa for the reduction of coliform bacteria and helminth eggs on vegetables. *Tropical Medicine and International Health*. 12: 40-50.
- Avazpoor M., Yousefipoor M.T., Dusty M., Mehdipour M., Seifipour F., Gholami Z. (2015). Determination of the level of parasitic infection (*Cryptosporidium* and *Giardia*) of the vegetables marketed in Ilam city. *Environmental Health Engineering and Management Journal*. 2: 33-40.
- Chalmers R.M., Katzer F. (2013). Looking for *Cryptosporidium*: the application of advances in detection and diagnosis. *Trends in Parasitology*. 29: 237-251.

- Checkley W., White A.C., Jaganath D., Arrowood M.J., Chalmers R.M., Chen X.M., Fayer R., Griffiths J.K., Guerrant R.L., Hedstrom L., Huston C.D. (2015). A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *The Lancet Infectious Diseases*. 15: 85-94.
- Damen J.G., Banwat E.B., Egah D.Z., Allanana J.A. (2007). Parasitic contamination of vegetables in Jos, Nigeria. *Annals of African Medicine*. 6: 115-118.
- Dixon B., Parrington L., Cook A., Pollari F., Farber J. (2013). Detection of *Cyclospora*, *Cryptosporidium*, and *Giardia* in ready-to-eat packaged leafy greens in Ontario, Canada. *Journal of Food Protection*. 76: 307-313.
- Duedu K.O., Yarnie E.A., Tetteh-Quarcoo P.B., Attah S.K., Donkor E.S., Ayeh-Kumi P.F. (2014). A comparative survey of the prevalence of human parasites found in fresh vegetables sold in supermarkets and open-aided markets in Accra, Ghana. *BMC Research Notes*. 7: 836.
- Fallah A.A., Pirali-Kheirabadi K., Shirvani F., Saei-Dehkordi S.S. (2012). Prevalence of parasitic contamination in vegetables used for raw consumption in Shahrekord, Iran: influence of season and washing procedure. *Food Control*. 25: 617-620.
- Gupta N., Khan D.K., Santra S.C. (2009). Prevalence of intestinal helminth eggs on vegetables grown in wastewater-irrigated areas of Titagarh, West Bengal, India. *Food Control*. 20: 942-945.
- Hadi A.M. (2011). Isolation and identification of intestinal parasites from vegetables from different markets of Iraq. *Bulletin of the Iraq Natural History Museum*. 11: 17-25.
- Insulander M., de Jong B., Svenungsson B. (2008). A food-borne outbreak of cryptosporidiosis among guests and staff at hotel restaurant in Stockholm County, Sweden. *Eurosurveillance*. 13: 1-2.
- Ma L., Sotiriadou I., Cai Q., Karanis G., Wang G., Wang G., Lu Y., Li X., Karanis P. (2014). Detection of *Cryptosporidium* and *Giardia* in agricultural and water environments in the Qinghai area of China by IFT and PCR. *Parasitology Research*. 113: 3177-3184.
- Maikai B.V., Baba-Onoja E.B.T., Elisha I.A. (2013). Contamination of raw vegetables with *Cryptosporidium* oocysts in markets within Zaria metropolis, Kaduna State, Nigeria. *Food Control*. 31: 45-48.
- Ortega Y.R., Roxas C.R., Gilman R.H., Miller N.J., Cabrera L., Taquiri C., Sterling C.R. (1997). Isolation of *Cryptosporidium parvum* and *Cyclospora cayetanensis* from vegetables collected in markets of an endemic region in Peru. *The American Journal of Tropical Medicine and Hygiene*. 57: 683-686.
- Petri W.A., Haque R., Lyster D., Vines R.R. (2000). Estimating the impact of amebiasis on health. *Parasitology Today*. 16: 320-321.
- Pires S.M., Vieira A.R., Perez E., Wong D.L.F., Hald T. (2012). Attributing human foodborne illness to food sources and water in Latin America and the Caribbean using data from outbreak investigations. *International Journal Food Microbiology*. 152: 129-138.
- Rahman J., Talukder A.I., Hossain F., Mahomud S., Islam A., Shamsuzzoha M. (2014). Detection of *Cryptosporidium* oocysts in commonly consumed fresh salad vegetables. *American Journal of Microbiological Research*. 2: 224-226.
- Ranjbar-Bahadori S.H., Mostoophi A., Shemshadi B. (2013). Study on *Cryptosporidium* contamination in vegetable farms around Tehran. *Tropical Biomedicine*. 30: 193-198.
- Rossle N.F., Latif B. (2013). Cryptosporidiosis as threatening health problem: a review. *Asian Pacific Journal of Tropical Biomedicine*. 3: 916-924.
- Rostami A., Ebrahimi M., Mehravar S., Omrani V.F., Fallahi S., Behniafar H. (2016). Contamination of commonly consumed raw vegetables with soil transmitted helminth eggs in Mazandaran province, Northern Iran. *International Journal of Food Microbiology*. 225: 54-58.
- Said D.E.S. (2012). Detection of parasites in commonly consumed raw vegetables. *Alexandria Journal of Medicine*. 48: 345-352.
- Sherbini G.T.E., Kamel N.O.H., Geneedy M.R., Temsah A.G. (2016). A comparative study of the occurrence of *Cryptosporidium parvum* oocysts found on fresh fruits and vegetables sold in supermarkets and open-aided markets. *International Journal of Current Microbiology and Applied Sciences*. 5: 760-768.
- Su G.L.S., Mariano C.M.R., Matti N.S.A., Ramos G.B. (2012). Assessing parasitic infestation of vegetables in selected markets in Metro Manila, Philippines. *Asian Pacific Journal of Tropical Diseases*. 2: 51-54.
- Tefera T., Biruksew A., Mekonnen Z., Eshetu T. (2014). Parasitic contamination of fruits and vegetables collected from selected local markets of Jimma Town, Southwest Ethiopia. *International Scholarly Research Notices*. 1: 1-7.