



Some Physicochemical Characteristics of Fish Products Sampled from Bulgarian Retail Markets

G. Zhelyazkov¹, D. Stratev^{*2}✉

1. Department of Biology and Aquaculture, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

2. Department of Food Hygiene and Control, Veterinary Legislation and Management, Faculty of Veterinary Medicine, Trakia University, 6000 Stara Zagora, Bulgaria

HIGHLIGHTS

- The highest level of water content and water activity was found in frozen fish products.
- The lowest pH value was found in the marinated fish products.
- Frozen fish products had the lowest salt and ash contents.
- Frozen Bulgarian fish products are probably more susceptible to spoilage than the other ones.

Article type

Original article

Keywords

Fish Products
Food Analysis
Food Safety
Bulgaria

Article history

Received: 1 Dec 2017
Revised: 25 Jan 2018
Accepted: 4 Feb 2018

Acronyms and abbreviations

a_w =Water activity

ABSTRACT

Background: The quality and safety of fish products is determined by chemical, physical, and microbiological parameters, important for satisfying consumers' requirements. Also, the freshness of fish is essential for evaluation of its quality. On the Bulgarian retail market, fish is commonly offered chilled, frozen, or processed. The purpose of this study was to determine some physicochemical characteristics of fish products sampled from Bulgarian retail markets.

Methods: During June to July 2017, this survey was performed on 45 samples from smoked, semi-dried, marinated, and frozen fish products sold in Bulgarian markets. The samples were collected from specialized stores for fish products, and transported to the laboratory for analysis. Water content, water activity (a_w), pH, salt content, and ash content were determined according to the standard protocols.

Results: The highest average water content was established in frozen products, followed by almost equal values in marinated, smoked, and semi-dried fish products. The average a_w value was also the highest in frozen products (0.975), it was almost the same in marinated and smoked (0.892); and the least in semi-dried fish products (0.905). Semi-dried (3.36%), marinated (3.19%), and smoked (3.03%) fish products had considerably higher average salt content than frozen ones (0.1%). The average pH value of marinated fish products (5.26) was lower than frozen (6.88), smoked (6.76), and also semi-dried (6.68) ones. Average ash content was substantially higher in smoked (7.16%), semi-dried (6.57%), and marinated (5.97%) fish products compared with frozen products (1.18%).

Conclusion: It is concluded that the frozen fish products sold in Bulgarian markets are probably more susceptible to spoilage than marinated, smoked, and semi-dried ones.

Introduction

The consumption of fish and fish products is beneficial for human health. Fish provides proteins, long-chain omega-3 polyunsaturated fatty acids, vitamins, and min-

erals (Borresen, 2008; Oguzhan and Angis, 2013).

The quality and safety of fish products is determined by chemical, physical, and microbiological parameters, important for satisfying consumers' requirements. Also, the freshness of fish is essential for evaluation of its quality

* Corresponding author. ✉ deyan.stratev@trakia-uni.bg

To cite: Zhelyazkov G., Stratev D. (2018). Some physicochemical characteristics of fish products sampled from Bulgarian retail markets. *Journal of Food Quality and Hazards Control*. 5: 33-36.

(Ababouch, 2006; Agustini et al., 2009; Alasalvar et al., 2011; Hall, 2012). On the Bulgarian retail market, fish is commonly offered chilled, frozen, or processed (smoked, semi-dried, marinated, and canned). The investigations on physicochemical parameters of fish products on the national market are few. Hence, the purpose of this study was to determine some physicochemical characteristics of fish products sampled from Bulgarian retail markets.

Materials and methods

During June to July 2017, this survey was performed on 45 samples from smoked, semi-dried, marinated, and frozen fish products sold in Bulgarian markets. Smoked fish comprised mackerel (*Scomber scombrus*; n=6), herring (*Clupea harengus membras*; n=6), and salmon fillet (*Salmo salar*; n=7). Semi-dried ones included mackerel (*Scomber scombrus*; n=6). Marinated fish samples were sprats (*Sprattus sprattus*; n=6) and Atlantic saury fillet (*Scomberesox saurus*; n=2). Frozen fish comprised

Alaska pollock fillet (*Theragra chalcogramma*; n=8) and pink salmon fillet (*Oncorhynchus gorbuscha*; n=4). The samples were collected from specialized stores for fish products, and transported to the laboratory for analysis.

Water content, water activity (a_w), pH, salt content, and ash content were determined according to the standard protocols described by AOAC (1990).

Results

The average water content in smoked, marinated, frozen, and semi-dried fish products were 60.34, 60.47, 80.75, and 41.80%, respectively; the average a_w values for the mentioned samples were 0.892, 0.892, 0.975, and 0.905, respectively. The mean pH values for smoked, marinated, frozen, and semi-dried products were determined as 6.76, 5.26, 6.88, and 6.68, respectively. The average salt and ash contents of the samples are shown in Figures 1 and 2. The details of physicochemical characteristics of fish products are presented in Table 1.




Figure 1: Average salt content (%) of different Bulgarian fish product samples




Figure 2: Average ash content (%) of different Bulgarian fish product samples

Table 1: Physicochemical characteristics of Bulgarian fish products samples

| Fish products | | Physicochemical characteristics (mean±SD) | | | | |
|---------------|------------------------|---|-------------|-----------|-----------|-----------------|
| | | Water content (%) | a_w | Salt (%) | pH | Ash content (%) |
| Smoked | Mackerel | 46.19±4.86 | 0.913±0.013 | 2.33±0.62 | 7.18±0.68 | 5.27±0.72 |
| | Herring | 67.75±2.02 | 0.880±0.004 | 4.61±0.19 | 7.19±0.08 | 11.03±0.74 |
| | Salmon fillets | 67.10±1.80 | 0.883±0.008 | 2.16±0.54 | 5.91±0.18 | 5.20±0.98 |
| Marinated | Sprat | 62.93±3.58 | 0.905±0.018 | 2.31±0.07 | 5.62±0.12 | 5.37±0.55 |
| | Atlantic saury fillets | 58.01±4.90 | 0.880±0.001 | 4.08±0.84 | 4.90±0.09 | 6.58±0.84 |
| Frozen | Alaska pollock fillets | 84.32±5.04 | 0.974±0.006 | 0.12±0.14 | 7.29±0.32 | 1.17±0.29 |
| | Pink salmon fillets | 77.19±2.18 | 0.977±0.001 | 0.09±0.06 | 6.48±0.08 | 1.19±0.11 |
| Semi-dried | Mackerel | 41.80±3.34 | 0.905±0.005 | 3.36±0.47 | 6.68±0.52 | 6.57±1.32 |

Discussion

In the present survey, the highest water content was established in frozen products, followed by marinated, smoked, and semi-dried fish products. The water in foods could be either bound or free (Syamaladevi et al., 2016). Free water determines the shelf life as it has an influence on enzymatic, chemical, and microbiological processes while bound water has no such effect (Agustini et al., 2009). Our results were in line with assumptions of Oguzhan and Angis (2013), that thermal processing decreased water content of fish products. Cardinal et al. (2004) reported average water content of 62.9% in smoked salmon, while Salán et al. (2006) found 68.01% water content in frozen salmon fillet. According to Adegunwa et al. (2013), the water content of smoked herring varied from 50.68 to 77.75%, corresponding with our data.

Similar to water content, a_w was in the highest level in frozen products (0.975), almost the same in marinated and smoked (0.892), and the least in semi-dried fish products (0.905). Agustini et al. (2009) reported a_w values between 0.57 and 0.87 in dried fish products sold in Indonesia. The a_w value is among the most important parameters during food preservation. It influences physical and chemical properties, the replication of microorganisms in foods and therefore, their shelf life. Most microbial enzymes are inhibited at a_w less than 0.85, and bacteria do not grow when a_w is less than 0.91 (Syamaladevi et al., 2016).

The absorption and distribution of salt within fish products in the view of Martinez et al. (2011) depend on the salt curing method, fish species, fillet thickness, fillet to salt ratio, muscle structure, and rigor mortis. On

the other hand, Bugueno et al. (2003) affirmed that salt curing prolongs shelf life of fish. In the present survey, salt concentrations of semi-dried (3.36%), marinated (3.19%), and smoked (3.03%) fish products had substantially higher salt content than frozen (0.1%) fish product samples. Similarly, Cardinal et al. (2004) reported average salt content of 3.1% in smoked salmon of European markets.

Marinating is among the oldest techniques for preservation of fish. Shelf life and safety of fish are due to added salt and organic acids, which reduce pH. It is proved that the pH value of 4.5 is sufficient to guarantee fish safety (Duyar and Eke, 2009). As expected, we found that the pH value of marinated fish products (5.26) was lower than that of frozen (6.88), smoked (6.76), and semi-dried ones (6.68). Anese and Gormley (1996) reported that the pH value of minced meat of cod and salmon was 6.8 and 6.6, respectively. Adegunwa et al. (2013) found that pH value in smoked herring was in range of 5.92-6.57 which was lower than the results of the present study.

In the current investigation, ash content was considerably higher in smoked (7.16%), semi-dried (6.57%), and marinated (5.97%) fish products compared with frozen (1.18%) fish products which were in accordance with the findings of Oguzhan and Angis (2013) and Salán et al. (2006). However, Adegunwa et al. (2013) established 10 times lower ash content in the smoked herring distributed in Nigeria. It has been announced that ash contents of seafood and fish products indicate that they are appropriate source of mineral substances like calcium, zinc, iron, as well as magnesium (Salindeho et al., 2014).

Conclusion

It is concluded that the frozen fish products sold in Bulgarian markets are probably more susceptible to spoilage than marinated, smoked, and semi-dried ones.

Conflicts of interest

There is no conflict of interest in this study.

Acknowledgments

This research was ethically approved by the local institutional review board. We thank Trakia University, 6000 Stara Zagora, Bulgaria for financial supports.

References

- Ababouch L. (2006). Assuring fish safety and quality in international fish trade. *Marine Pollution Bulletin*. 53: 561-568.
- Adegunwa M.O., Adebawale A.A., Olisa Z.G., Bakare H.A. (2013). Chemical and microbiological qualities of smoked herring (*Sardinella eba*, Valenciennes 1847) in Odeda Ogun State, Nigeria. *International Journal of Microbiology Research and Reviews*. 1: 85-87.
- Agustini T.W., Darmanto Y.S., Susanto E. (2009). Physicochemical properties of some dried fish products in Indonesia. *Journal of Coastal Development*. 12: 73-80.
- Alasalvar C., Miyashita K., Shahidi F., Wanasundara U. (2011). Handbook of seafood quality, safety and health applications. John Wiley and Sons, UK.
- Anese M., Gormley T.R. (1996). Effects of dairy ingredients on some chemical, physico-chemical and functional properties of minced fish during freezing and frozen storage. *LWT-Food Science and Technology*. 29: 151-157.
- Association of Official Analytical Chemists (AOAC). (1990). Official methods of analyses. 15th edition. Association of official analytical chemists, Washington DC.
- Borresen T. (2008). Improving seafood products for the consumer. Elsevier, Boca Raton.
- Bugueno G., Escriche I., Martínez-Navarrete N., del Mar Camacho M., Chiralt A. (2003). Influence of storage conditions on some physical and chemical properties of smoked salmon (*Salmo salar*) processed by vacuum impregnation techniques. *Food Chemistry*. 81: 85-90.
- Cardinal M., Gunnlaugsdottir H., Bjoernevik M., Ouisse A., Vallet J.L., Leroi F. (2004). Sensory characteristics of cold-smoked Atlantic salmon (*Salmo salar*) from European market and relationships with chemical, physical and microbiological measurements. *Food Research International*. 37: 181-193.
- Duyar H.A., Eke E. (2009). Production and quality determination of marinade from different fish species. *Journal of Animal and Veterinary Advances*. 8: 270-275.
- Hall G.M. (2012). Fish processing technology. Springer Science and Business Media, London.
- Martinez O., Salmerón J., Guillén M.D., Casas C. (2011). Characteristics of dry- and brine-salted salmon later treated with liquid smoke flavouring. *Agricultural and Food Science*. 20: 217-227.
- Oguzhan P., Angis S. (2013). Effects of processing methods on the sensory, mineral matter and proximate composition of rainbow trout (*Oncorhynchus mykiss*) fillets. *African Journal of Food Science and Technology*. 4: 71-75.
- Salán E.O., Galvão J.A., Oetterer M. (2006). Use of smoking to add value to the salmoned trout. *Brazilian Archives of Biology and Technology*. 49: 57-62.
- Salindeho N., Purnomo H., Yuniarta, Kekenusa J. (2014). Physicochemical characteristics and fatty acid profile of smoked skipjack Tuna (*Katsuwonus pelamis*) using coconut fiber, nutmeg shell and their combination as smoke sources. *International Journal of ChemTech Research*. 6: 3841-3846.
- Syamaladevi R.M., Tang J., Villa-Rojas R., Sablani S., Carter B., Campbell G. (2016). Influence of water activity on thermal resistance of microorganisms in low-moisture foods: a review. *Comprehensive Reviews in Food Science and Food Safety*. 15: 353-370.